

PHENOTYPIC CHARACTERIZATION OF DONKEYS IN BENISHANGUL GUMUZ NATIONAL REGIONAL STATE

Amine MUSTEFA^{1✉}, Abraham ASSEFA¹, Manaye MISGANAW¹, Fasil GETACHEW², Solomon ABEGA³, Abebe HAILU¹ and Yibrehu EMSHAW¹

¹Ethiopian Biodiversity Institute, P. O. Box 30726, Addis Ababa, Ethiopia

²ILRI (International Livestock Research Institute), P. O. Box 5689, Addis Ababa, Ethiopia

³EIAR (Ethiopian Institute of Agricultural Research, Debre-Zeit Agricultural Research Center), P. O. Box Bishoftu, Ethiopia

✉Email: aminemustefa32@gmail.com;  ORCID: 0000-0003-3788-4156

➤Supporting Information

ABSTRACT: Fifteen morphometric measurements and eighteen qualitative traits were recorded on 323 randomly sampled adult donkeys (123 jacks and 200 jennets) to meet an objective of characterizing a heterogeneous donkey population of three phenotypic types (Sinnar, the locals and their crosses) found in Benishangul Gumuz region of Ethiopia. The General Linear Model and non-parametric test (chi-square) procedures of SAS software were used for the analysis of the morphometric data and qualitative traits, respectively. Means were separated using the Tukey-Kramer test. The studied morphometric measurements were significantly affected by the phenotypic type and partially affected by sex and sample location /district. Sinnar donkeys were significantly bigger and heavier than the local and the crosses. However, for some of the morphometric measurements no significance difference was observed between Sinnar and crosses implying the presence of heterosis. Heart girth measurements for Sinnar, local and crosses were 110.61 ± 0.436 , 106.18 ± 0.448 and 108.87 ± 1.251 , respectively. Body weight estimates of 127.26 ± 1.277 , 113.40 ± 1.312 and 121.13 ± 3.665 kg were obtained for Sinnar, the local and the crosses. There was sexual size dimorphism and depending on the type of morphometric trait either jacks or jennets show significantly ($P < 0.05$) higher values as compared to the opposite sex. Jacks had wider chest (22.61 vs 22.09 cm.), thicker (24.40 vs 23.24 cm.), and longer cannon bone (31.00 vs 31.32 cm) than jennets, while the jennets possess wider hip (33.00 vs 31.69 cm.) and longer body (90.49 vs 88.52 cm.), back (67.37 vs 66.17), ear (24.42 vs 23.90 cm) and heavier estimated weight (122.47 vs 118.71 kg) than the jacks. Limited location effect was recorded showing donkeys from Guba district were comparatively the largest. Majority of the studied donkeys possess white abdominal color, unpigmented hoof and muzzle, short and medium hair size, plain body color pattern with long dorsal stripe without leg stripe, straight face and sloppy rump profile, medium tail length and thickness at the base of the tail. Qualitative differences ($P < 0.05$) were also observed among the class categories. Further molecular level studies could supplement the current study and provide more refined classification of the various genotypes in the studied area. Similarly, characterization of the local donkeys found in other parts of the region and the Abyssinian donkeys in the adjoining areas is required.

Keywords: Heart girth, Morphometric, Qualitative, Phenotypic types, Sinnar

INTRODUCTION

Ethiopia is endowed with diverse domestic, aquatic and wild animal genetic resources. The diverse ecology Ethiopia has and its position as a route of entry to domestic animals from Asia to Africa has resulted in the presence of diverse animal genetic resources. Despite the presence of the resources adequate characterization work is lacking and the information on the state of the animal genetic resources is incomplete to support sustainable utilization and conservation of the resources (EBI, 2016). This is more so for the equine genetic resources of the country.

Donkeys, like other livestock species, have an important place in rural and urban communities in Ethiopia. Even if donkeys have not been serving as a food source to humans in Ethiopia due to religious and cultural taboos, they make significant economic contribution in all the regions. Donkeys specifically are important for transport of goods in urban, peri-urban and rural areas. In the latter case they also serve in transporting humans, threshing cereal crops and plowing of land pairing with oxen.

The donkey (*Equus asinus*) is indigenous to the African continent and its wild progenitor is usually considered to be the Nubian wild ass (Blench, 2000). Ethiopia possesses the largest donkey population in the world with 8,439,220 donkeys (CSA, 2017; FAO, 2015). Even if, donkey is the least studied species in the country, previously four types of donkeys were recognized; namely Jimma, Abyssinian, Ogaden and Sinnar based on their phenotypic and physical characteristics like average size and coat colour (Befikadu et al., 2015). However, more recent nationwide study identified six distinct domestic donkey populations namely Abyssinian, Afar, Hararghe, Ogaden, Omo and Sinnar (Kefena 2012, Kefena 2014; EBI 2016). The same study showed that, most of the variations in the parameters of morphological

RESEARCH ARTICLE
 DOI: <https://dx.doi.org/10.36380/sci.2020.ojaf14>
 PII: S222877012000004-10
 Received: November 13, 2019
 Revised: December 27, 2019

characteristics vary with eco-geographical patterns and biophysical resources. Therefore, other morphometric variables and corporal indices need to be further incorporated and used to fully characterize and describe donkey populations in Ethiopia (Befikadu et al., 2015). Like all the other regions found within the country, Benishangul Gumuz region possess different donkey phenotypes and their crosses including the country's largest donkey type (Sinnar). Hence this study was aimed at revealing the phenotypic characteristics (qualitative and quantitative parameters) of the different donkey breeds found in the region to be used as input for a further conservation and sustainable utilization of the resources

MATERIALS AND METHODS

Description of the study areas

The study was conducted in Benishangul Gumuz Regional State. Benishangul Gumuz Regional State is one of the nine regional states which is located in the western part of Ethiopia between longitude 34° 10'N and 37° 40'E; and latitude 09° 17'N and 12° 06' N. It shares border with Amhara Region in the north and north east, South Sudan in the west, Gambella Region in the south, Oromia Region in the south east. The total area of the Region is approximately 50,380 square kilometers with altitude ranging from 580 to 2731 meters above sea level. About 75% of the Region is low land, 24% is semi-high land and 1% is high land. The capital city of the Region is Asossa, located at a distance of 659 kms west of Addis Ababa (Chekol and Getnet, 2010).




Figure 1 - Map of the study areas

Sampling technique and sample size

The studied animals (adult donkeys for phenotypic characterization; morphometric and qualitative records) were sampled randomly from five districts (Guba, Menge, Wenbera, Sirba Abay and Kurmuk) within the region. Heterogeneous donkey populations of three phenotypic types (Sinnar, the local donkey and their crossbreds) were used for the qualitative records and morphometric measurements. A total of 323 full-mouthed adult donkeys (123 jacks and 200 jennets) composed of 157 Sinnar, 148 local and 18 crossbred donkeys were measured for linear and circular morphometric traits and described for the qualitative traits.

Measurement and data collection

The sample size determination and identification of traits for morphometric measurements and qualitative description were based on FAO guideline (FAO, 2012). Fifteen quantitative/ morphometric measurements (Heart girth, Height at wither, Height at back, Height at rump, Body length, Back length, Neck length, Head length, Canon bone length, Fore leg length, Hip width, Chest width, Chest depth, Canon circumference and Ear length) and 18 qualitative characteristics (coat hair size, body color pattern, body (coat) color, abdominal color, head color, ear tip color, tail switch color, hoof color, and muzzle color, ear shape, dorsal stripe, leg stripe, shoulder stripe, face profile, back profile, rump

profile, tail length and thickness at the base of the tail) were recorded from each individual.

Donkeys were carefully handled by trained laborers and made to stand squarely on flat grounds. Morphometric measurements and qualitative data recording were made by separate individuals. According to a study by [Košťuková \(2015\)](#), who reported that the growth of donkeys terminates after the age of 5 years and all donkeys in the study were past this age. Body weight was estimated from the above measurements by the formula from [Pearson and Ouassat \(2000\)](#).

$$\text{Live weight (kg)} = (\text{heart girth [cm]} \times 2.12) \times (\text{body length [cm]} \times 0.688) / 3801$$

Table 1 - Sampled number of animals by sex and by breed and proportion of each breeds.

Breed /genotype	Jacks	Jennets	Total	Proportion
Sinnar	57	100	157	0.48
Local	58	90	148	0.46
Crosses	8	10	18	0.06
Total	123	200	323	1.00
Proportion	0.38	0.62	1.00	-

Statistical analysis

Data entry and management were done using Microsoft Excel. Analysis of data on quantitative measurements was carried out using the GLM procedure of SAS 9.0 software. Means were compared using Tukey-Kramer ([SAS, 2002](#)). Similarly, analysis of qualitative traits was carried out using the non-parametric test (chi-square) procedure of SAS 9.0 software. The model used for the analysis of quantitative data: $Y_{ijk} = \mu + A_i + B_j + C_k + e_{ijk}$, where Y_{ijk} is an observation, μ is the overall mean, A_i is the fixed effect of sex, B_j is the fixed effect of the breed group, C_k is the fixed effect of district and e_{ijk} is the random error attributed to the n th observation. Interaction effects were found to be non-significant in most cases and were removed from the analysis model.

RESULTS

Morphometric measurements

The overall mean, standard error (SE), minimum and maximum value, and coefficients of variation (CV) of the collected morphometric measurements are presented in [table 2](#). For all morphometric traits measured the coefficient of variation was within the range of 5.11 and 8.27%. Relatively higher coefficient of variation (13.11%) was calculated for estimated body weight implying higher variation in terms of body weight. The difference between the minimum and maximum value is sizeable in most cases. A range of 40 cm for height at wither, 37 cm for body length, and a range of about 104 kg for body weight were observed.

Table 2 - Overall least square mean , SE, CV, Minimum and Maximum of body measurements of the donkey populations in Benishangul Gumuz region.

Traits	Overall Mean ± SE	Minimum	Maximum	CV (%)
Hearth girth (cm)	108.7 ± 0.31	93.0	128.0	5.18
Height at wither (cm)	100.2 ± .030	79.0	119.0	5.31
Height at back (cm)	102.0 ± 0.31	89.0	122.0	5.41
Height at rump (cm)	102.3 ± 0.30	90.0	121.0	5.22
Body length (cm)	89.9 ± 0.30	68.0	105.0	6.00
Back length (cm)	67.3 ± 0.24	57.0	86.0	6.34
Neck length (cm)	49.2 ± 0.23	38.0	60.0	8.27
Head length (cm)	42.9 ± 0.12	36.0	50.0	5.11
Canon bone length (cm)	31.4 ± 0.12	26.0	39.0	6.92
Fore leg length (cm)	68.3 ± 0.26	54.0	86.0	6.92
Hip width (cm)	32.6 ± 0.11	27.0	40.0	6.60
Chest width (cm)	22.3 ± 0.09	18.0	29.0	7.63
Chest depth (cm)	49.1 ± 0.15	42.0	59.0	5.65
Canon circumference (cm)	23.7 ± 0.09	20.0	29.0	6.70
Ear length (cm)	23.9 ± 0.10	19.0	28.0	7.27
Body weight (kg)	121.2 ± 0.93	77.8	181.6	13.82

SE=Standard error, SD=Standard deviation, CV=Coefficient of variation and cm=Centimeter.

Table 3 - Least square means (cm) and pairwise comparison of body measurements with standard error in each breeds/populations: 1) jacks.

Traits	Breed groups/ phenotypic types			p value
	Sinnar	Local	Cross	
N	57	58	8	
Heart girth	110.4±0.72 ^a	104.9±0.71 ^b	107.9±1.89 ^{ab}	***
Height at wither	104.0±0.61 ^a	96.5±0.60 ^b	101.4±1.60 ^a	***
Height at back	106.5±0.65 ^a	98.8±0.64 ^b	103.1±1.70 ^a	***
Height at rump	106.2±0.67 ^a	98.9±0.66 ^b	102.3±1.74 ^{ab}	***
Body length	90.9±0.70 ^a	86.2±0.69 ^b	86.9±1.82 ^{ab}	***
Back length	67.9±0.52 ^a	64.7±0.51 ^b	65.9±1.36 ^{ab}	***
Neck length	50.4±0.53 ^a	46.9±0.53 ^b	50.1±1.39 ^{ab}	***
Head length	43.9±0.27 ^a	42.2±0.26 ^b	43.7±0.69 ^{ab}	***
Canon bone length	33.0±0.30 ^a	30.8±0.30 ^b	32.3±0.78 ^{ab}	***
Fore leg length	70.7±0.59 ^a	66.4±0.58 ^b	68.1±1.55 ^{ab}	***
Hip width	32.0±0.26	31.4±0.26	31.0±0.68	NS
Chest width	23.0±0.21 ^a	22.1±0.20 ^b	22.0±0.53 ^{ab}	**
Chest depth	50.1±0.33 ^a	47.2±0.33 ^b	49.9±0.86 ^a	***
Canon circumference	24.8±0.21 ^a	23.9±0.20 ^b	24.6±0.54 ^{ab}	*
Ear length	24.0±0.25 ^a	22.9±0.24 ^b	25.8±0.64 ^a	**
Body weight	126.3±2.14 ^a	108.9±2.10 ^b	116.4±5.57 ^{ab}	***

N= Number of observations, *p<0.05, **p<0.01, ***p<0.0001, NS=Not Significant

In jacks, with the exception of hip width, donkey phenotypic type had a significant ($P<0.01$) effect on the measured traits and it is indicated that almost all body measurements were highest for Sinnar jacks (Table 3) followed by the crossbred type. In almost all cases the difference between the Sinnar and the crossbred jacks is not significant while the difference between the Sinnar and the local is significant in all cases.

The difference between the local and the crosses reached significance level only for height at wither and at back, chest depth and ear length. The Sinnar and crossbred jacks have larger chest depth and taller height at withers than the local jacks. Positive heterosis of more than one percent was calculated for height at withers, neck length, head length, canon bone length, chest depth and ear length. The highest heterosis of close to ten per cent was observed for ear length.

In jennets, significant ($P<0.05$) difference was observed between phenotypic types for all morphometric traits (Table 4). In all cases Sinnar jennets along with the crosses have significantly higher values than the local donkeys. No significant difference was observed between Sinnar jennets and the crossbreds. Despite the fact that the crosses would of various types (F1, F2, F3, back cross etc.) and blood levels heterosis of more than one per cent was calculated for body length, fore leg length, hip width, chest width, chest depth, ear length and body weight of jennets.

Table 4 - Least square means (cm) and pairwise comparison of body measurements with standard error in each breeds/populations: 2) jennets.

Traits	Breed groups/ phenotypic types			p value
	Sinnar	Local	Cross	
N	100	90	10	
Heart girth	110.8±0.53 ^a	107.1±0.56 ^b	109.8±1.66 ^{ab}	***
Height at wither	102.6±0.6 ^a	97.4±0.49 ^b	100.7±1.43 ^{ab}	***
Height at back	104.3±0.46 ^a	98.9±0.49 ^b	101.3±1.43 ^{ab}	***
Height at rump	104.7±0.44 ^a	99.6±0.47 ^b	102.4±1.39 ^{ab}	***
Body length	92.4±0.50 ^a	89.7±0.53 ^b	92.0±1.56 ^{ab}	**
Back length	69.0±0.42 ^a	66.8±0.45 ^b	66.2±1.32 ^{ab}	**
Neck length	50.7±0.39 ^a	48.6±0.42 ^b	48.9±1.22 ^{ab}	**
Head length	43.7±0.20 ^a	41.8±0.21 ^b	43.2±0.62 ^{ab}	***
Canon bone length	32.1±0.18 ^a	30.4±0.20 ^b	31.4±0.58 ^{ab}	***
Fore leg length	70.1±0.42 ^a	66.8±0.45 ^b	69.2±1.33 ^{ab}	***
Hip width	33.6±0.21 ^a	32.5±0.23 ^b	33.4±0.67 ^{ab}	**
Chest width	22.6±0.17 ^a	21.6±0.19 ^b	22.6±0.54 ^{ab}	**
Chest depth	50.2±0.26 ^a	48.2±0.28 ^b	50.1±0.82 ^{ab}	***
Canon circumference	23.6±0.15 ^a	23.0±0.16 ^b	22.9±0.47 ^{ab}	*
Ear length	24.6±0.16 ^a	23.7±0.17 ^b	24.4±0.50 ^{ab}	**
Body weight	128.6±1.54 ^a	117.1±1.64 ^b	125.5±4.83 ^{ab}	***

N= number of observations, *p<0.05, **p<0.01, ***p<0.0001, NS=Not Significant

The combined data analysis has revealed that there is significant difference between the phenotypic types for all traits and Sinnar donkeys were superior in all cases but the differences between Sinnar and the crosses in some cases was not significant (Table 5). Sinnar donkeys and their crosses have significantly higher ($P<0.05$) chest depth, ear length, canon bone length and height at withers. The three types were significantly different from each other for height at back and rump. Positive heterosis as calculated from the combined data was found to be more than one per cent for head length, canon bone length, chest depth and ear length. The highest (4.22%) was calculated for ear length.

Comparison of jacks and jennets have shown that for some of the morphometric variables there is sexual size dimorphism where either the jack or jennet could show higher values as compared to the other sex (Table 6). Jacks have shown significantly ($P<0.05$) higher values for height at back, cannon bone length, chest width and cannon bone circumference while jennets have significantly higher values for body length, back length, hip width, ear length and body weight. There was no significant ($P>0.05$) difference between the sexes for heart girth, height at withers, height at rump, neck length, head length, fore leg length and chest depth.

The donkeys were sampled from five locations /districts to consider if there are environmental differences; however, location effects were limited to few of the recorded traits. Body length, back length, neck length, head length, fore leg length and ear length are the traits which have shown significant ($P<0.05$) fluctuations as the sampling location /district differs. Based on this, foreleg length of donkeys from Wenbera district and head length of donkeys from Sirba Abay district were the shortest among the donkey populations from the other districts. More or less, the results show measurements of donkey populations from Guba district were comparably the largest showing above mean performances.

Table 5 - Least square means (cm) and pairwise comparison of body measurements with standard error in each breeds/populations: 3) all sexes.

Traits	Breed groups/ phenotypic types			p value
	Sinnar	Local	Cross	
N	157	148	18	
Heart girth	110.6 ± 0.44 ^a	106.2 ± 0.45 ^b	108.9 ± 1.25 ^{ab}	***
Height at wither	103.2 ± 0.37 ^a	97.1 ± 0.38 ^b	101.0 ± 1.07 ^a	***
Height at back	105.3 ± 0.38 ^a	98.9 ± 0.39 ^c	102.2 ± 1.10 ^b	***
Height at rump	105.3 ± 0.38 ^a	99.4 ± 0.39 ^c	102.4 ± 1.09 ^b	***
Body length	91.5 ± 0.41 ^a	88.0 ± 0.43 ^b	89.7 ± 1.19 ^{ab}	***
Back length	68.5 ± 0.33 ^a	65.9 ± 0.34 ^b	65.9 ± 0.96 ^b	***
Neck length	50.5 ± 0.32 ^a	47.8 ± 0.33 ^b	49.4 ± 0.91 ^{ab}	***
Head length	43.9 ± 0.16 ^a	42.0 ± 0.16 ^b	43.5 ± 0.46 ^a	***
Canon bone length	32.5 ± 0.16 ^a	30.6 ± 0.17 ^b	31.9 ± 0.47 ^a	***
Fore leg length	70.3 ± 0.35 ^a	66.7 ± 0.36 ^b	68.8 ± 1.04 ^{ab}	***
Hip width	32.8 ± 0.17 ^a	31.9 ± 0.17 ^b	32.3 ± 0.48 ^{ab}	**
Chest width	22.8 ± 0.14 ^a	21.9 ± 0.14 ^b	22.4 ± 0.39 ^{ab}	***
Chest depth	50.2 ± 0.21 ^a	47.8 ± 0.22 ^b	50.0 ± 0.60 ^a	***
Canon circumference	24.2 ± 0.12 ^a	23.5 ± 0.13 ^b	23.7 ± 0.35 ^{ab}	**
Ear length	24.3 ± 0.14 ^a	23.4 ± 0.14 ^b	24.8 ± 0.39 ^a	***
Body weight	127.3 ± 1.28 ^a	113.4 ± 1.31 ^b	121.1 ± 3.67 ^{ab}	***

N= Number of observations, * $p<0.05$, ** $p<0.01$, *** $p<0.0001$

Table 6 - Least square means and pairwise comparison of body measurements with standard error in each sexes.

Body variables	Sex		p value
	Jacks	Jennets	
N	123	200	
Heart girth	108.2±0.59	108.9±0.53	NS
Height at wither	100.7±0.51	100.1±0.46	NS
Height at back	102.9±0.52	101.4±0.47	**
Height at rump	102.7±0.51	102.1±0.46	NS
Body length	88.5±0.56	90.9±0.51	***
Back length	66.2±0.45	67.4±0.41	**
Neck length	48.9±0.43	49.6±0.39	NS
Head length	43.3±0.22	42.9±0.20	NS
Canon bone length	32.0±0.22	31.3±0.20	**
Fore leg length	68.7±0.48	68.5±0.43	NS
Hip width	31.7±0.23	33.0±0.21	***
Chest width	22.6±0.18	22.1±0.17	**
Chest depth	49.2±0.28	49.5±0.26	NS
Canon circumference	24.4±0.17	23.2±0.15	***
Ear length	23.9±0.19	24.4±0.17	**
Body weight	118.7±1.73	122.5±1.56	*

N= Number of observations, * $p<0.05$, ** $p<0.01$, *** $p<0.0001$, NS=Not Significant

Table 7 - Least square means (cm) and pairwise comparison of body measurements with standard error in each districts.

Traits	Sampled Location /District					p value
	Guba	Menge	Wenbera	Sirba Abay	KurmuK	
N	58	84	89	59	33	
Heart girth	108.7±0.81	108.8±0.64	108.5±0.68	109.7±0.77	107.1±0.97	NS
Height at wither	100.2±0.69	100.4±0.55	99.9±0.58	101.8±0.66	99.8±0.83	NS
Height at back	101.9±0.71	101.7±0.56	101.9±0.59	102.8±0.68	102.4±0.86	NS
Height at rump	102.7±0.70	102.1±0.56	101.8±0.59	103.1±0.67	102.1±0.85	NS
Body length	89.0±0.77 ^{ab}	89.8±0.61 ^{ab}	88.5±0.64 ^b	91.3±0.73 ^a	90.0±0.93 ^{ab}	*
Back length	66.8±0.62 ^{ab}	66.4±0.49 ^b	66.4±0.52 ^b	68.4±0.59 ^a	65.9±0.74 ^b	*
Neck length	49.8±0.59 ^{ab}	48.1±0.47 ^b	49.0±0.49 ^{ab}	49.9±0.56 ^a	49.3±0.71 ^{ab}	*
Head length	43.3±0.30 ^a	43.6±0.24 ^a	43.0±0.25 ^a	41.9±0.28 ^b	43.7±0.36 ^a	***
Canon bone length	31.8±0.30	31.2±0.24	31.5±0.25	31.7±0.29	32.1±0.36	NS
Fore leg length	69.5±0.65 ^a	69.1±0.52 ^a	66.1±0.55 ^b	69.2±0.62 ^a	69.1±0.79 ^a	***
Hip width	32.5±0.31	32.6±0.25	32.5±0.26	31.6±0.30	32.5±0.37	NS
Chest width	22.7±0.25	22.5±0.20	21.9±0.21	22.3±0.24	22.3±0.30	NS
Chest depth	49.0±0.39	49.9±0.31	49.2±0.32	48.9±0.37	49.5±0.47	NS
Canon circumference	23.7±0.23	23.9±0.18	23.8±0.19	23.7±0.22	23.9±0.27	NS
Ear length	24.8±0.25 ^a	23.8±0.20 ^b	23.9±0.21 ^b	24.3±0.24 ^{ab}	23.9±0.31 ^{ab}	**
Body weight	120.2±2.48	121.1±1.88	119.4±1.98	124.8±2.25	117.5±2.85	NS

N= Number of observations, *p<0.05, **p<0.01, ***p<0.0001, NS=Not Significant

Qualitative characteristics

The qualitative characteristics of the studied donkey population under the effect of breed, sex and source of the animals is presented in Table 8 and 9. Genotype and environmental factors like sex, animal source and district affects the qualitative characteristics of the studied donkeys. The donkey populations studied possess the following qualitative characteristics; white abdominal color, unpigmented hoof and muzzle, short and medium hair coat cover, plain body color pattern with long dorsal stripe (leg stripe in some cases), straight face and sloppy rump profile, medium tail length and thickness at the base of the tail. The studied qualitative characteristics were affected by different genetic and environmental factors. Breed group has a significant effect on the studied traits except on coat description, abdominal color, face profile and tail length while sex affects few of the qualitative characteristics recorded like coat description, back profile and tail thickness at the base. Similarly source of the animal affects most of the traits except body color pattern, ear shape, leg strip and rump profile.

The results showed dorsal body color of the studied donkey populations was affected ($P<0.05$) by breed group of the donkeys. Based on this result, dorsal body color of Sinnar donkeys were 18% brown, 17% dark brown, 15% white, 13% light red, 8% dark gray, 8% gray dun and 21% others. While the locals were 24% gray dun, 23% bay dun, 16% dark gray dun, 13% gray/roan, 9% brown and 15% others. Similarly, the crossbreds were 22% bay dun, 22% dark gray dun, 17% light red, 11% black, 11% dark brown and 17% other colors. The head color of the Sinnar donkeys was 31% white, 11% brown, 10% light red, 10% dark brown, 9% black and 29% others. However, the head color of the local donkeys was 21% gray dun, 18% bay dun, 16% dark gray dun, 14% gray/roan, 10% brown and 21% others. The crossbreds were also 28% dark gray dun, 17% bay dun, 17% white, 11% light red, 11% dark brown, 11% gray/roan and 5% others in head color. The ear tips color of Sinnar donkeys was 47% dark brown, 21% black, 10% brown, 9% light red and 13% others, while the local donkeys were 63% dark brown, 28% black and 9% other colors. Similarly, the crossbreds were 44% dark brown, 39% black and 17% other in colors of their ear tips. The dominant tail switch color of the studied donkeys was black, with dark brown; almost all (89%) of the crossbreds possess black tail switch color, similarly most (81%) of the locals tail switch color was black with 16% dark brown. The results also showed that 60% of the Sinnar donkeys had black tail switch color with 20% dark brown and 20% other colors.

The study revealed that local and cross donkeys possess plain body color pattern while some of the Sinnar donkeys were shaded. Almost 70 percent of the Sinnar donkey breeds ear shape was round edged while most of the local and crossbreds ear was straight edged in shape. Almost all local (93%) and crossbreds (83%) possess long dorsal stripe while this feature was observed on half of the Sinnar donkey breeds. 40 % of the local donkeys had leg stipe, however, it was not seen on most of the Sinnar and its crossbred donkeys. Most local donkeys and the crossbreds had either short or long shoulder stripe, while, it was absent on more than half of the Sinnar donkeys. Almost 50% of the Sinnar donkeys back profile was convex, however, in most of the crossbreds and local donkeys it was hollow.

The results also showed that most of the short coat hair size was possessed by jacks while the jennets had medium coat description. Most of the jacks back profile was hollow while the jennets were hollow. Most of the jacks had small to medium thickness at the base of their tail, however it was medium to large for the jennets. Purchased animals had short coat hair size while the born ones had medium coat description.

There was no dorsal stripe on half of the purchased donkeys while half of them possess long. On the other hand, most of the donkeys born on-farm possess long dorsal stripe. Similarly, there was no shoulder stripe on more than half (52%) of the purchased donkeys while there was either short or long shoulder stripe for the donkeys born there. Most of the purchased donkeys back profile was convex while half of the borne ones possess hollow back profile. Short and medium tail length was observed on the donkeys born there while the purchased donkeys tail length was medium to long.

Animals were sampled from different locations/districts to have a representative picture of the study area (Benishangul Gumuz region). Most of the qualitative characteristics of the studied donkey populations were significantly ($P<0.05$) influenced by location /district. Based on this, ear shape, leg stripe, shoulder stripe, back profile, rump profile, tail length and thickness at the base of the tail are some of the traits which show significant differences among districts. The results show most of the donkeys from Kurmuk district had round edged ear shape while straight edged ear shape was seen on most of the donkeys sampled from Menge district. Most of the donkey populations from the sampled locations do not possess leg stripe while 50% of the donkeys from Guba and Kurmuk districts had a leg stripe.

The results also revealed that most of the donkeys from Sirba Abay district had a hollow back profile while straight back profile was observed on above half of the donkeys from Guba district. On the other hand, above half of the donkeys from Menge district had convex back profile.

Table 8 - Percentage of qualitative traits in each breed/populations, sexes and animal sources.

Traits	Breeds/populations			Sex		Animal source	
	Sinnar	Local	Cross	Jacks	Jennets	Born	Purchased
N	157	148	18	123	200	232	91
Coat hair size (Chi-square)		NS			***		**
Short	48	39	39	63	31	38	57
Medium	47	53	61	36	60	55	38
Long	5	8	0	1	9	7	5
Body color pattern (Chi-square)		***			NS		NS
Plain	75	100	94	89	87	90	81
Shaded	25	0	6	11	13	10	19
Ear shape (Chi-square)		***			NS		NS
Rounded	69	22	28	48	43	42	53
Straight	31	78	72	52	57	58	47
Dorsal strip (Chi-square)		***			NS		***
Absent	50	7	17	29	27	22	45
Long	49	93	83	70	72	77	55
Short	1	0	0	1	1	1	0
Leg strip (Chi-square)		**			NS		NS
Present	19	39	17	29	27	30	22
Absent	81	61	83	71	73	70	78
Shoulder stripe (Chi-square)		***			NS		***
Absent	52	9	17	30	31	22	52
Long	21	46	44	32	35	39	21
Short	27	45	39	38	34	39	27
Face profile (Chi-square)		NS			NS		*
Convex	27	38	33	36	31	37	22
Straight	72	61	67	63	68	62	78
Concave	1	1	0	1	1	1	0
Back profile (Chi-square)		**			*		**
Hollow	32	47	61	32	45	47	24
Straight	19	23	0	20	20	20	19
Convex	49	30	39	48	35	33	57
Ramp profile (Chi-square)		**			NS		NS
Flat	22	36	39	37	25	33	22
Sloppy	65	59	61	56	65	59	68
Roofy	13	5	0	7	10	8	10
Tail length (Chi-square)		NS			*		*
Short	25	30	33	26	30	32	18
Medium	50	40	22	37	48	43	46
Long	25	30	45	37	22	25	36
Tail base thickness (Chi-square)		*			***		*
Narrow	33	17	17	34	19	20	36
Medium	44	56	55	59	45	53	44
Wide	23	27	28	7	36	27	20

N= number of observations, * $p<0.05$, ** $p<0.01$, *** $p<0.0001$, NS=Not Significant

Table 9 - Percentage of qualitative traits in each districts.

Traits	Location /district					Chi-square
	Guba	Menge	Wenbera	Sirba Abay	Kurmuk	
N	58	84	89	59	33	
Coat hair size						NS
Short	53	39	43	44	39	
Medium	40	54	50	51	61	
Long	7	7	7	5	0	
Body color pattern						NS
Plain	84	90	89	83	91	
Shaded	16	10	11	17	9	
Ear shape						***
Rounded	43	26	52	47	73	
Straight	57	74	48	53	27	
Dorsal strip						NS
Absent	26	32	34	17	27	
Long	74	68	64	83	73	
Short	0	0	2	0	0	
Leg strip						***
Present	43	27	11	31	42	
Absent	57	73	89	69	58	
Shoulder stripe						**
Absent	33	35	35	15	30	
Long	22	39	25	56	27	
Short	45	26	40	29	42	
Face profile						**
Convex	26	27	48	20	39	
Straight	74	70	52	80	61	
Concave	0	3	0	0	0	
Back profile						***
Hollow	16	30	40	76	45	
Straight	55	13	24	0	0	
Convex	29	57	36	24	56	
Ramp profile						**
Flat	21	32	27	42	24	
Sloppy	77	55	66	54	55	
Roofy	2	13	7	4	21	
Tail length						***
Short	21	11	29	44	55	
Medium	45	55	38	37	39	
Long	34	34	33	19	6	
Tail base thickness						***
Narrow	31	36	21	3	33	
Medium	48	44	59	49	49	
Wide	21	20	20	48	18	

N= number of observations, *p<0.05, **p<0.01, ***p<0.0001, NS=Not Significant

DISCUSSION

Effect of breed group

The results revealed that there were differences among the studied breeds/populations indicating Sinnar donkeys were significantly bigger than the local donkeys while some similarities were observed with the crosses which might be due to heterosis effect. Heterosis of various magnitude was calculated for the various traits studied. However, as the crosses are of diverse type (F1, F2, back cross etc.) and blood level it will be difficult to interpret the heterosis effect obtained in this study in both sexes. The big body size, height and length in the measured traits of Sinnar indicated that the breed is highly adaptable to the hot environment with good reproduction ability (Beja-Pereira et al., 2004; Marshall 2007; Rossel et al., 2008). These results are in line with the results of Kefena et al., (2011) and Tsega and Lemma (2015) who reported Sinnar donkeys are the tallest of all donkey populations in Ethiopia and they are also an excellent desert adapted animals used for riding and breeding. Therefore, due to these special characteristics of the Sinnar donkey, Kefena et al. (2011) has reported that its ancestral trunk might be different from the rest of the donkey breeds/populations but this needs to be supported by further genetic studies. Some of the characteristics of desert adapted donkeys include fairly bigger body sizes and similarities in coat color patterns among donkey populations of arid and semi-arid lowlands (Beja-Pereira et al., 2004; Marshall, 2007; Rossel et al., 2008). Differences in breed groups in this study were in line with the results of Kosťuková et al. (2015). The possible cause of differences among the studied breed groups might be due to the differences in domesticated ancestors, ecology and biophysical resources for the overall body build and local environment and history (Beja-Pereira et al., 2004; Kefena et al., 2014; Gubitiz et al., 2000). The local breeds were smaller and lighter than the Sinnar donkeys which is suitable for different purposes which is in line with the

results of [Sargentini et al. \(2018\)](#) who reported that the Amiata donkeys' biometrics (small-medium sized) was suitable for different purposes.

The dominant body colors of Sinnar donkeys from this study were white and light colors with short to medium hair while, most of the locals and crossbreds possess gray dun and bay dun with medium hair size. These results are in line with the results of study by [Tsega and Lemma \(2015\)](#) in Gondar, Ethiopia, who reported Sinnar donkeys possess white body color with short hair and crossing them with Abyssinian donkeys often results in hairier, longer, saggy and dark colored hybrids.

Effect of sex

Eight of the fifteen morphometric measurements were significantly affected by sex; jacks were dominant over the jennets in four traits (height at back, chest width, canon bone length and canon circumference) while the jennets were dominant over jacks in five traits (body length, back length, ear length, hip width and body weight). These results were in line with the results of [Košťuková et al., \(2015\)](#) and partially with the results of [Folch and Jordana \(1997\)](#) on Catalonian donkeys. The magnitude of sexual dimorphism in the current study is quite higher than what has been reported for Catalonian donkeys where only Eight out of twenty-six morphometric measurements showed significant differences between sexes ([Folch and Jordana, 1997](#)). The report of [Koubek \(1933\)](#) also showed that male donkeys have lower heart girth and stronger shin than the jennets. Jacks in this study were stronger (wide chest and thick canon) than the jennets which is in line with the results of [Andersson \(1994 cit. in Purzyc et al., 2007\)](#) who reported that, there are specific physiological and biochemical processes in jacks, which results in them being stronger than the jennets. In line with the current study the chest circumference and hip width in female individuals is mainly influenced by physiological processes during gestation and by metabolic traits that differ from those in males ([Koubek, 1933](#)). Similarly, height at wither and at rump were not significantly different among the sexes while the jennets had shorter height at back than the jacks which might be due to the fetus load which pull their belly down during the pregnancy period. On the other hand, it might also be due to their loss of strength to hold up the load they carry in comparison to jacks.

Effect of location

Location affects only six traits (body, back, head, neck, ear and fore leg length) out of the eighteen morphometric measurements. In the study by [Tsega and Lemma \(2015\)](#) on Sinnar donkeys in Gondar, Ethiopia, differences in morphometric measurements and qualitative characteristics were recorded among different sampling location due to differences in ecological selection regimes, history or both. However, such huge significant differences were not recorded in the present study which may be due to the closeness of the locations sampled. However also, the donkeys from the locations/ districts which are on the border with South Sudan shows some dominance over the others as these districts may serve as the gate ways of the Sinnar donkeys to the region. Most of the qualitative characteristics of the purchased donkeys approaches the characteristics of Sinnar donkeys, showing most of the farmers were purchasing Sinnar donkeys preferring their heaviness. The white color and short hair size of the Sinnar donkeys might help them in physiological adaptation of the hot environments.



Local Male



Sinnar Male



Cross Male



Local Female



Sinnar Female



Cross Female

CONCLUSION

The donkeys in Benishangul Gumuz region were characterized based on FAO guidelines. Accordingly, three phenotypic types were covered; the Sinnar, the locals and their crossbreds. Significant differences were recorded among the three breed groups on the studied morphometric measurements and qualitative characteristics. Based on these results the Sinnar donkeys were the tallest, widest and strongest with some similarities with their crosses. Some sex effect was observed on the overall performances while, its effect was limited on the interaction effect with the breed groups. However, jacks and jennets were dominating each other on equal basis. Jacks had wide chest, thick canon and long height at rump and cannon bone, while the jennets possess wide hip and long body, back, neck and ear. Significant differences were observed between sampled location for some of the studied traits. Qualitative differences were also recorded among the studied class categories. Further molecular level studies are required to characterize the differences among the studied donkey breed groups. Similarly, characterization of the local donkeys found in other parts of the region and the Abyssinian donkeys in adjoining areas is required.

DECLARATIONS

Corresponding Author

E-mail: aminemustefa32@gmail.com; ORCID: 0000-0003-3788-4156

Acknowledgements

The authors are highly indebted to the animal owners as without their permission the completion of this work would not have been possible. Our special appreciation also goes to the head and staff of Ethiopian Biodiversity Institute (especially Animal Biodiversity Directorate), and the district agricultural offices found in the study area who actively participated during the data collection process.

Authors' Contribution

AM contribute on data analysis and the write up of the manuscript, AA, MM, FG conceived the study and collect data, SA review the manuscript, AH and YE contribute on entering data. All authors read and approved the final manuscript.

Funding

This research was funded by the Ethiopian biodiversity institute.

Conflict of Interests

The authors have not declared any conflict of interests.

REFERENCES

- Befikadu Z, Kiflay W and Sanjoy KP (2015) Conservation of Indigenous Donkey Breeds of Ethiopia: A Review. *International Journal of Interdisciplinary and Multidisciplinary Studies*, 2 (6): 13-22.
- Beja-Pereira A, England P R, Ferrand N, Jordan S, Bakhiet A O, Abdalla M A, Mashkour M, Jordana J, Taberlet P and Luikart G (2004) African origin of domestic donkey. *Science*, 304: 1781.
- Blench RM (2000) A history of donkeys, wild asses and mules in Africa. In: Blench, Roger M., MacDonald, Kevin C. (Eds.), *The origin and development of African livestock: archaeology, genetics, linguistics and ethnography*. UCL Press, pp. 339–354.
- Chekol K and Getnet A (2010) Public Finance Review: Benishangul-Gumuz Regional State Pp. 83
- CSA (Central Statistical Agency) (2017) Agricultural sample survey 2016/2017. Report on livestock and livestock characteristics. Addis Ababa, Ethiopia. pp. 188
- EBI (Ethiopian Biodiversity Institute) (2016) Ethiopian National Strategy and Plan of Action for conservation and utilization of Animal Genetic Resources, Addis Ababa, Ethiopia. Pp. 114
- FAO (Food and Agricultural Organization of the United Nations) (2015) In: Scherf B. & Pilling D. (Eds.), *The second report on the state of world's animal genetic resources for food and agriculture*. FAO, Rome, Italy.
- FAO (Food and Agricultural Organization of the United Nations) (2012) Phenotypic characterization of animal genetic resources. Food and Agriculture Organization of United Nations. Rome, Italia. *Animal Production and Health Guidelines* 11, 144.
- Folch P and Jordana J (1997) Characterization, reference ranges and the influence of gender on morphological parameters of the endangered Catalanian donkey breed. *Journal of Equine Veterinary Science*, 17 (2): 102–111.
- Gubitz T, Thorpe R S and Malhotra A (2000) Phylo-geographic and natural selection in the Tenerife gecko *Tarentola delalandii*: testing historical and adaptive hypothesis. *Molecular Ecology*, 9: 1213–1221.
- Kefena E, Beja-Pereira A, Han JL, Haile A, Mohammed YK and Dessie T (2011) Eco-geographical structuring and morphological diversities in Ethiopian donkey populations. *Livestock Science* 141, 232–241 <http://dx.doi.org/10.1016/j.livsci.2011.06.011>

- Kefena E, Dessie T, Tegegne A, Beja-Pereira A, Yusuf Kurtu M, Rosenbom S and Han JL (2014) Genetic diversity and matrilineal genetic signature of native Ethiopian donkeys (*Equus asinus*) inferred from mitochondrial DNA sequence polymorphism. *Livestock Science*, <http://dx.doi.org/10.1016/j.livsci.2014.06.006>
- Kefena E (2012) Equine genetic resources of Ethiopia. PhD. dissertation Haramaya University, Dire Dawa. Ethiopia.
- Košťuková M, Černohorská H, Bihunčová I, Oravcová I, Sobotková E and Jiskrová I (2015) Characteristics of morphological parameters of donkeys in the Czech Republic. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 63 (2): 419–424 <http://dx.doi.org/10.11118/actaun201563020419>
- Koubek K (1933) Speciální zootechnika – Chov koní. Praha: Státní zemědělské nakladatelství Praha.
- Marshall F. (2007) African pastoral perspectives on domestication of the donkey: a first synthesis. In: Denham, T., Iriarte, J., Vrydaghs, L. (Eds.), *Rethinking Agriculture: Archaeological and Ethnoarchaeological Perspectives*. Left Coast Press, Walnut Creek, California, USA.
- Pearson R A and Ouassat M (2000) A Guide to Body Condition Scoring and Live Weight Estimation of Donkeys. Centre for Tropical Veterinary Medicine, University of Edinburgh, p. 21
- Purzyc H, Kobryn H, Komosa M and Bojarski J (2007) Ocena eksterieru konia huculskiego na podstawie wybranych wskaźników morfometrycznych (część I). *Acta Scientiarum Polonorum – Medicina Veterinaria*, 6: 47–64.
- Rossel S, Marshall F, Peters J, Pilgram T, Adams M D and O'Connor D. (2008) Domestication of the donkey: timing, processes and indicators. *Proceedings of the National Academy of Sciences of the United States of America*, 105: 3715–3720.
- Sargentini C, Tocci R, Martini A and Bozzi R (2018) Morphological characterization of Amiata donkey through Multivariate analyses. *Revista Brasileira de Zootecnia*, 47: e20170310, <https://doi.org/10.1590/rbz4720170310>
- SAS (Statistical Analysis System). (2002) Institute Inc., Cary, NC, USA.
- Tsega A and Lemma A (2015) Phenotypic characteristics and sexual behavior of Sennar Jacks (*Equus asinus*). *Ethiopian Veterinary Journal*, 19 (1): 11 – 21.