



Effects of Dietary Inclusion of Maggot (*Hermetia illucens*) and Corn on Productivity of Starter-Stage Native Chickens

Sabrang Panuntun¹, Cahya Utama^{2*}, and Bambang Sulistiyanto³

Faculty of Animal and Agricultural Science, Diponegoro University, Semarang, Indonesia

*Corresponding author's Email: cahyasetyautama@gmail.com

Received: March 06, 2025, Revised: April 10, 2025, Accepted: May 12, 2025, Published: June 25, 2025

ABSTRACT

Native chickens require innovative feeding strategies to enhance productivity. This study aimed to examine the dietary effects of using maggot and corn meal (MCM) on the performance and feed digestibility of native chickens. The study used a complete randomized design, with a total of 200 one-day-old native chicks randomly allocated into four groups, each with five replicates. The first group received 100% basal feed (T0). The second group received 95% basal feed and 5% MCM (T1), the third group received 90% basal feed and 10% MCM (T2), and the fourth group received 85% basal feed and 15% MCM (T3). Group T1 exhibited significantly higher live weight, daily body weight gain, crude protein digestibility, crude fiber digestibility, and crude fat digestibility compared to other treatments. The T3 group showed the highest feed intake, whereas the best feed conversion ratio (FCR) was observed in the T0 group. It is concluded that a 5% inclusion of MCM in the feed has beneficial effects on growth performance and digestibility of native chickens.

Keywords: Black soldier fly, Combination feed, Corn, Digestibility, Native chicken, Performance

INTRODUCTION

According to the BPS-National Statistics (2023), the total population of native chickens in Indonesia has recently been increasing, serving as an important alternative protein source. The population of native chickens in Indonesia rose from 300.1 million in 2018 to 308.6 million in 2022 (BPS, 2023). Despite their significance, native chickens have exhibited relatively slow growth and high feed conversion values (Promket and Ruangwittayanusorn, 2021). Feed quality is the main factor that supports the growth of native chickens. Low feed quality prevents native chickens to grow and develop optimally (Jachimowicz et al., 2022). The high price of manufactured feed encourages farmers to seek affordable yet nutritionally adequate alternatives (Alshelmani et al., 2021). Developing innovative feeds using the advantages of locally available ingredients presents a viable solution. As an energy source, maize can be combined with protein sources to meet nutritional sufficiency (Barszcz et al., 2022). Black Soldier Fly (BSF) larvae (*Hermetia illucens*), also known as maggots, are a promising protein source, containing 44.26% protein content and equipped with

various types of essential amino acids (Schiavone et al., 2019). Compared to soybean meal, maggot exhibits superior amino acid content, including histidine (2.91% vs. 1.23%), leucine (6.79% vs. 3.65%), lysine (8.25% vs. 2.88%), methionine (1.84% vs. 0.62%), threonine (3.86% vs. 1.84%), and valine (5.54% vs. 2.3%) (Surendra et al., 2020; Aguirre et al., 2024). However, the Black soldier fly maggot has a limiting factor in chitin content derived from the exoskeleton (Purkayastha and Sarkar, 2020). High-pressure heating is expected to loosen the bonds in chitin and bring out the resistant starch of maize. Chitin will then look like a surface with large pores and dense piles of microfibrils. After steaming, the surface of chitin becomes smooth without microfibrils, and the pores increase in depth and spread over the entire surface (Tan et al., 2015). Resistant starch can be utilized by intestinal microflora as a food substrate during the digestive process, so it has the potential to become a prebiotic (Qin et al., 2020).

The current study aimed to evaluate the dietary effects of BSF maggot and corn, processed by high-pressure heating, on the productivity of native chicks. This study utilizes maize as an energy source and BSF maggot as a protein source, combined via high-pressure heating.

Final weight

The final body weight measurement was carried out by weighing via a digital scale at the end of the rearing period (on the 28th day).

Daily body weight gain

Body weight was measured using digital scales. Measurement of Daily Body Weight Gain (DBWG) in g/chicken/day was calculated following Formula 1 (Utama et al., 2020):

$$DBWG = \frac{\text{Final weight} - \text{initial weight}}{28 \text{ (days of observation)}} \quad (\text{Formula 1})$$

Feed intake

Feed intake data were collected daily by calculating the difference between the given feed and the residual feed. Weighing of residual feed was done before feeding in the morning.

Feed conversion ratio

Feed Conversion Ratio (FCR) was calculated based on the ratio between the amount of ration consumed and the measured weight gain.

$$FCR = \frac{\text{Amount of feed consumed (g)}}{\text{Body weight gain (g)}} \quad (\text{Formula 2})$$

Digestibility values

Measurement of digestibility values was performed on 28-day-old chicks by sampling 2 chickens per replicate. Total collection methods were conducted according to Utama et al. (2020). Each replicate group of chickens was kept in a battery cage (2 chickens/repetition/cage), and 6 chickens were placed in each cage to get endogenous excreta. Chickens were moved to the battery cage and allowed to adjust to the conditions for 12 hours by providing feed and water. The transfer of chickens to battery cages was done at night and under low temperatures to minimize stress on the chickens. Afterwards, all treated and endogenous chickens were fasted for 24 hours to eliminate feed residues in the digestive tract. Chickens were fed 100 g of treatment feed, and the excreta were collected for 48 hours. Chickens for endogenous excreta collection were kept fasted and given water *ad libitum*. Endogenous chicken excreta were collected for 44 hours. Excreta were sprayed once every hour with 1N HCl to capture and reduce nitrogen evaporation. The collected excreta were cleaned of contaminants, then dried by natural drying for 2 days at

28-34°C. During the drying process, the excreta were covered using an insect net to prevent flies and other contaminants from entering, and were still routinely sprayed with HCl. The samples were then ground and put into plastic clips according to the replication to be tested for crude protein, crude fiber, crude fat, and nitrogen content.

Crude Protein Digestibility (CPD), Crude Fiber Digestibility (CFD), and Crude Fat Digestibility/Ether Extract Digestibility (EED) were calculated following Formulas 3-5, respectively (Moningkey et al., 2019). The nitrogen retention (NR) was calculated using Formula 6 (Sibbald, 1980):

$$\text{CPD (\%)} = \frac{(\text{CP consumption} - \text{CP excreta})}{\text{CP consumption}} \times 100\% \quad (\text{Formula 3})$$

$$\text{CFD (\%)} = \frac{(\text{CF consumption} - \text{CF excreta})}{\text{CF consumption}} \times 100\% \quad (\text{Formula 4})$$

$$\text{EED (\%)} = \frac{(\text{EE consumption} - \text{EE excreta})}{\text{EE consumption}} \times 100\% \quad (\text{Formula 5})$$

$$\text{NR (\%)} = \frac{(\text{Fd} \times \text{Nf}) - ([\text{E} \times \text{Ne}] - [\text{En} \times \text{Nen}])}{(\text{Fd} \times \text{Nf})} \times 100\% \quad (\text{Formula 6})$$

where Fd is the feed consumed (g), Nf is the nitrogen feed (%), Ne is the nitrogen excreta (%), E is the total excreta (g), En is the total excreta endogenous (g), and Nen is the nitrogen endogenous (%).

Data analysis

Data were analyzed using SPSS, version 23.0. The data were tested for normality and homogeneity and were analyzed using analysis of variance (ANOVA). Significant differences ($p < 0.05$) identified via one-way ANOVA were further analyzed using Duncan's Multiple Range Test (DMRT).

RESULT AND DISCUSSION

Growth performance

The dietary inclusion of maggot-corn meal (MCM) significantly influenced ($p < 0.05$) the final weight, daily body weight gain (DBWG), feed intake (FI), and feed conversion ratio (FCR) of starter-stage native chickens (Table 2).

Table 1. Effects of different dietary MCM levels on final weight, daily body weight gain, feed intake, and feed conversion in the starter-stage of native chickens

Parameter	Treatment	T0	T1	T2	T3
Final weight (g/chicken) *		374.7 ± 26.14 ^{ab}	384.6 ± 4.32 ^a	367.6 ± 7.72 ^{ab}	359.3 ± 18.6 ^b
Daily body weight gain (g/chicken/day) *		12.71 ± 0.94 ^{ab}	12.34 ± 0.16 ^a	12.06 ± 0.28 ^{ab}	11.8 ± 0.65 ^b
Feed intake (g/chicken/day) *		21.21 ± 0.32 ^d	24.38 ± 0.17 ^c	25.51 ± 0.18 ^b	26.1 ± 0.12 ^a
Feed conversion*		1.59 ± 0.01 ^a	1.78 ± 0.03 ^b	1.95 ± 0.04 ^c	2.04 ± 0.1 ^c

Source: Primary research data (2024); *Different superscript letters in each row show significant differences ($p < 0.05$). The first group received 100% basal feed (T0). The second group received 95% basal feed and 5% MCM (T1), the third group received 90% basal feed and 10% MCM (T2), and the fourth group received 85% basal feed and 15% MCM (T3).

Final weight and daily body weight gain

The various levels of MCM in the diets had a significant effect ($p < 0.05$) on the final weight and daily body weight gains of starter-stage native chicks. The highest final weight and DBWG values were found in the second group (T1) (384.6 ± 4.32 g/chicken and 12.34 ± 0.16 g/chicken/day) whereas the lowest final weight and DBWG indexes were found in the fourth treatment group (T3) (359.3 ± 18.6 g/chicken and 11.8 ± 0.65 g/chicken/day). This suggests that MCM is effective as an alternative feed only up to a 5% inclusion level. Factors that can affect final weight are body weight gain and feed conversion. Utama et al. (2024) noted that the final weight is related to both feed conversion and body weight gain, as the more feed nutrients are converted into tissues that form organs, which can increase body weight gain, so that the final weight increases. Maggot and corn meal levels of 10 and 15% in the diet decreased the final weight. This can be attributed to the chitin content in MCM. Kastalani et al. (2021) stated that chitin can inhibit protein digestion because it can form complex bonds with proteins and bind nitrogen in amino acids. According to Ramdani et al. (2018), chickens need protein for metabolism in the meat formation process. Warisman et al. (2024) also stated that the 5.4% of chitin in steamed maggot decreased the digestibility value and nutritional parameters of the chicken ration. A decreased level of digestibility will affect the feed conversion.

Feed intake

The effect of different levels of MCM on the diet had a significant effect ($p < 0.05$) on the feed intake of starter-

stage native chicken. The addition of MCM to the diet tends to increase the ration consumption of native chickens. Feed intake increased with MCM inclusion, peaking in T3 (26.1 g/head/day) and then T2 (25.51 g/head/day) and T1 (24.38 g/head/day) compared to the control (21.21 g/head/day). It is assumed that the chitin content is increased by increasing the content of MCM in the diet. According to Daniar and Herdyastuti (2019), chitin is grouped among the animal fibers whose structure is similar to cellulose, with a crude fiber content of 72.52%. An increase in crude fiber will trigger the rate of digesta in the intestine so that feed quickly leaves the digestive tract, the livestock feed hungry, and continue to consume rations (Jha and Misshra, 2021). Chitin will increase the digesta rate, so that the chickens will quickly feel hungry and continue to consume feed.

Feed conversion ratio (FCR)

The feed conversion ratio of starter-stage native chickens was also influenced by various MCM levels. The control group (T0) achieved the best FCR (1.59 ± 0.01), while T3 had the poorest (2.04 ± 0.1). The usage of MCM in the ration tends to increase FCR. This can be attributed to the values of nitrogen retention. Abun et al. (2023) stated that higher nitrogen retention results in more efficient feed conversion. Utama et al. (2024) added that high nitrogen retention values indicate high quality feed protein containing balanced essential and non-essential amino acids, which are easily absorbed and converted into body weight. This is relevant to the nitrogen retention value in Table 3, showing that higher nitrogen retention values lead to higher FCR values.

Table 3. Effect of different dietary MCM levels on crude protein, crude fiber, crude fat digestibility, and nitrogen retention in the starter-stage of native chickens

Parameter	Treatment	T0	T1	T2	T3
Crude protein digestibility*		80.73 ± 0.69 ^b	83.35 ± 0.9 ^a	81.75 ± 0.8 ^b	78.27 ± 1.11 ^c
Crude fiber digestibility*		45.72 ± 1.82 ^b	53.01 ± 1.49 ^a	47.6 ± 0.85 ^b	39.64 ± 1.55 ^c
Crude fat digestibility*		81.78 ± 0.95 ^c	92.33 ± 1.35 ^a	91.5 ± 0.69 ^a	85.48 ± 1.39 ^b
Nitrogen Retention*		88.10 ± 1.38 ^a	88.75 ± 0.4 ^a	83.26 ± 0.68 ^b	80.71 ± 2.75 ^b

Source: Primary research data (2024); *Different superscript letters in each row show significant differences ($p < 0.05$). The first group received 100% basal feed (T0). The second group received 95% basal feed and 5% MCM (T1), the third group received 90% basal feed and 10% MCM (T2), and the fourth group received 85% basal feed and 15% MCM (T3).

Feed digestibility

The findings revealed that there were significant effects ($p < 0.05$) arising from the type of treatment on crude protein digestibility, crude fiber digestibility, crude fat digestibility, and retention of nitrogen, as presented in Table 3.

Crude protein digestibility

Dietary MCM inclusion significantly affected ($p < 0.05$) Crude Protein Digestibility (CPD) in starter-stage native chicks. with the highest CPD observed in T1 ($83.35 \pm 0.9\%$) and the lowest in T3 ($78.27 \pm 1.11\%$). This shows that MCM is effective up to 5%, and the higher levels will reduce protein digestibility. In comparison, increasing the composition of MCM and reducing the composition of soybean meal in the diet also affected protein digestibility. This is thought to be due to the chitin content in MCM. Chitin can form complex bonds with proteins and bind amino acid nitrogen (Kastalani et al., 2021). A study by Warisman et al. (2024) showed that the steamed maggot decreased the digestibility values and the nutritional parameters of the chicken ration due to the chitin content. Sánchez-Muros et al. (2014) stated that poultry lack chitinase enzymes, limiting their ability to absorb chitin.

Crude fiber digestibility

Different MCM levels in the diet had a significant effect ($p < 0.05$) on Crude Fiber Digestibility (CFD) among starter-stage native chicks. As presented in Table 3, the highest CFD was observed in the second group (T1) at 53.01%, with the third group (T2) at 47.6%, the first group (T0) at 45.72%, and finally reaching at 39.64% in the fourth group (T3) as the lowest CFD. This finding shows that the addition of MCM at 10% and 15% levels tends to reduce the level of crude fiber digestibility in native chickens. It seems the reason is the fact that the chitin content at 10% and 15% MCM level causes an increase in digesta rate, so that the process of digesting crude fiber by intestinal microbes only lasts for a short time. Chitin in feed will accelerate the rate of digesta; the faster the digesta rate is, the shorter the digestion process will be (Has et al., 2013; Nirwana et al., 2021). Morgan et al. (2022) noted that an extremely short digestion rate leads to a short time available for digestive enzymes to degrade nutrients, which can reduce feed digestibility.

Crude fat digestibility

Varying MCM levels in diets significantly affected ($p < 0.05$) Crude Fat Digestibility (CFD) in starter-stage

native chicks. The highest CFD was obtained in the second group (T1) at 92.33%, followed by the third group (T2) at 91.5%, and then the fourth group (T3) at 85.48%, finally achieving 81.78% as the lowest in the first group (T0). Using MCM in the ration effectively increased crude fiber digestibility up to 10%, but it decreased CDF at a 15% level. The MCM feeding significantly increased crude fat digestibility compared to the control treatment ($p < 0.05$). This is probably because maggot has a type of fat that is easily digested. Black soldier fly larvae contain 37.34% monounsaturated fatty acids and 20.45% polyunsaturated fatty acids and are dominated by medium-chain fatty acids (C8-C12) (Kim and Rhee, 2016; Alifian et al., 2019). Unsaturated fatty acids can also affect the performance of lipase enzymes to increase fat absorption (Adhami et al., 2021).

Nitrogen retention

Nitrogen retention (NR) was highest in T1 (88.75%) and lowest in T3 (80.71%). According to Sibbald and Wolynetz (1985), a factor that helps increase nitrogen retention value is the production of quality feed, which is easily converted into weight. This indicates that the use of MCM at the 5% level can improve feed quality, seen from the nitrogen retention value perspective. The decrease in nitrogen retention value in T2 and T3 is thought to be due to the chitin content in the MCM. The addition of MCM results in an increase in the chitin content in the feed. Kastalani et al. (2021) stated that chitin can bind nitrogen from amino acids, reducing the digestibility value of protein. Belluco et al. (2013) and Lee et al. (2022) argue that chitin causes a decrease in the digestibility of feed ingredients and can bind the nutrients needed.

CONCLUSION

It is suggested that maggot and corn meal be replaced with soybean meal up to 5% to increase live weight, daily weight gain, and digestibility values of crude protein, crude fiber, and crude fat of native chickens. Further studies on changes in the micronutrient structure of MJS are needed to determine its potential as a functional feed ingredient.

DECLARATIONS

Acknowledgements

We express our gratitude to Daud, Tazkia, Bilqis, Eka, Anis, Fatma, and Reisha for their invaluable assistance in the research activities.

Funding

This research was self-funded without external financial assistance.

Author's contributions

Sabrang Panuntun conducted the research, collected and analyzed the data, and drafted the manuscript. Cahya Utama and Bambang Sulistiyanto reviewed and edited the manuscript. All authors have read and approved the final version of the manuscript.

Competing interests

The authors have declared that no competing interests exist.

Ethical considerations

Plagiarism, consent to publish, misconduct, data fabrication and/or falsification, double publication and/or submission, and redundancy have been checked by all the authors.

Availability of data and materials

All data generated during the study are included in this article. Any additional information is available upon reasonable request.

REFERENCES

- Abun A, Putra RRO, and Haetami K (2023). Metabolizable energy and nitrogen retention of local chickens fed rations containing fermented catfish waste. *GSC Biological and Pharmaceutical Sciences*, 25(3): 138-148. DOI: <https://www.doi.org/10.30574/gscbps.2023.25.3.0531>
- Adhami BAK, Amirkolaei H, Oraji M, Kazemifard, and Mahjoub S (2021). Effects of lysophospholipid on rainbow trout (*Oncorhynchus mykiss*) growth, biochemical indices, nutrient digestibility and liver histomorphometry when fed fat powder diet. *Aquaculture Nutrition*, 27: 1779-1788. DOI: <https://www.doi.org/10.1111/anu.13315>
- Aguirre L, Cámara L, Smith A, Fondevila G, and Mateos GG (2024). Apparent metabolizable energy and ileal amino acid digestibility of commercial soybean meals of different origins in broilers. *Poultry Science*, 103(7): 103786. DOI: <https://www.doi.org/10.1016/j.psj.2024.103786>
- Alifian MD, Sholikin MM, Evvyernie D, and Nahrowi (2019). Potential fatty acid composition of *H. illucens* oil reared on different substrates. *IOP Conference Series: Materials Science and Engineering*, 546: 0620021-5. DOI: <https://www.doi.org/10.1088/1757-899X/546/6/062002>
- Alshelmani MI, Abdalla EA, Kaka U, and Basit MA (2021). Nontraditional feedstuffs as an alternative in poultry feed. *Advances in poultry nutrition research*. IntechOpen., London, United Kingdom, pp. 1-214. DOI: <http://www.doi.org/10.5772/intechopen.95946>
- Barszcz M, Anna T, and Marcin T (2022). Poultry nutrition. *Physical Sciences Reviews*, 9(2): 611-650. DOI: <https://www.doi.org/10.1515/psr-2021-0122>
- Belluco S, Losasso C, Maggioletti M, Alonzi CC, Paoletti MG, and Ricci A (2013). Edible insects in a food safety and nutritional perspective: A critical review. *Comprehensive Reviews in Food Science and Food Safety*, 12(3): 296-313. DOI: <https://www.doi.org/10.1111/1541-4337.12014>
- BPS-National statistics (2023). Native chicken population by province. National standardization agency. Available at: <https://www.bps.go.id/statistics-table/2/NDc2IzI=/populasi-ayam-buras-menurut-provinsi.html>
- Has H, Yunianto VD, and Sukanto B (2013). The effectivity of fermented mulberry leaves with rumen liquor as broiler feed on final body weight, dry matter and crude fiber digestibility, and metabolic energy. *Animal Production*, 15(3): 173-179. Available at: <http://www.animalproduction.net/index.php/JAP/article/view/431>
- Herdyastuti N and Daniar AV (2019). The effect of chitin size and soaking time on decreasing cholesterol levels on quail eggs. *National Seminar on Chemistry 2019 (SNK-19)*. Atlantis Press, pp. 91-94. DOI: <https://www.doi.org/10.2991/snk-19.2019.1>
- Kastalani K, Kusuma ME, Herlinae H, and Yemima Y (2021). Effect of addition of maggot and rice bran-based feed-in basal feed against live weight of broiler carcass and giblet. *Ziraa'ah*, 46(1): 44-52. DOI: <https://www.doi.org/10.31602/zmip.v46i1.3732>
- Jachimowicz, K, Winiarska-Mieczan A, and Tomaszewska E (2022). The impact of herbal additives for poultry feed on the fatty acid profile of meat. *Animals*, 12(9): 1054. DOI: <https://www.doi.org/10.3390/ani12091054>
- Jha R and Mishra P (2021). Dietary fiber in poultry nutrition and their effects on nutrient utilization, performance, gut health, and on the environment: A review. *Journal of Animal Science and Biotechnology*, 12: 51. DOI: <https://www.doi.org/10.1186/s40104-021-00576-0>
- Kim SA and Rhee MS (2016). Highly enhanced bactericidal effects of medium chain fatty acids (caprylic, capric, and lauric acid) combined with edible plant essential oils (carvacrol, eugenol, β -resorcylic acid, trans-cinnamaldehyde, thymol, and vanillin) against *Escherichia coli* O157: H7. *Food Control*, 60: 447-454. DOI: <https://www.doi.org/10.1016/j.foodcont.2015.08.022>
- Lee JH, Kim TK, Cha JY, Jang HW, Yong HI, and Choi YS (2022). How to develop strategies to use insects as animal feed: Digestibility, functionality, safety, and regulation. *Journal of Animal Science and Technology*, 64(3): 409-431. DOI: <https://www.doi.org/10.5187/jast.2022.e27>
- Moningkey AF, Wolayan FR, Rahasia CA, and Regar MN (2019). Digestibility of the organic matter, crude fiber and crude fat in broilers ration containing pumpkin waste meal (*Cucurbita moschata*). *Zootec*, 39(2): 257-265. DOI: <https://www.doi.org/10.35792/zot.39.2.2019.24870>
- Morgan N, Bhuiyan MM, and Hopcroft R (2022). Non-starch polysaccharide degradation in the gastrointestinal tract of broiler chickens fed commercial-type diets supplemented with either a single dose of xylanase, a double dose of xylanase, or a cocktail of non-starch polysaccharide-degrading enzymes. *Poultry Science*, 101(6): 101846. DOI: <https://www.doi.org/10.1016/j.psj.2022.101846>
- Nirwana, Mujnisa A, and Jamilah (2021). Length and weight of small intestine and digestion rate of quail, with the addition of beluntas leaf flour (*Pluchea indica* L.) to the ration. *IOP Conference Series: Earth and Environmental Science*, 788(1): 012073. DOI: <https://www.doi.org/10.1088/1755-1315/788/1/012073>
- Nuningtyas YF, Aulanni'Am AA, Sjoftan O, Nurgartiningih VMA, Furqon A, Lestari SP, and Natsir MH (2024). Dietary protein levels in the small intestine and carcass traits of cross-breed chickens. *Journal of World's Poultry Research*, 14(1): 88-97. DOI: <https://www.doi.org/10.36380/jwpr.2024.9>

- Promket D and Ruangwittayanusorn K (2021). The comparatives of growth and carcass performance of the Thai native chicken between economic selection (Chee KKU12) and natural selection (Chee N). *Veterinary Integrative Sciences*, 19(2): 247-257. DOI: <https://www.doi.org/10.12982/VIS.2021.022>
- Purkayastha D and Sarkar S (2020). Physicochemical structure analysis of chitin extracted from pupa exuviae and dead imago of wild black soldier fly (*Hermetia illucens*). *Journal of Polymers and the Environment*, 28(2): 445-457. DOI: <https://www.doi.org/10.1007/s10924-019-01620-x>
- Qin S, Zhang K, Applegate TJ, Ding X, Bai S, Luo Y, Wang J, Peng H, Su Z, Xuan Y et al. (2020). Dietary administration of resistant starch improved caecal barrier function by enhancing intestinal morphology and modulating microbiota composition in meat duck. *British Journal of Nutrition*, 123(2): 172-181. DOI: <https://www.doi.org/10.1017/S0007114519002319>
- Ramdani OP, Suthama N, and Atmomarsono U (2018). The effect of protein levels and L-lysine rations of fatty Kampong chicken in 12 weeks. *Journal of Indonesian Veterinary Science*, 13(4): 388-394. DOI: <https://www.doi.org/10.31186/jspi.id.13.4.388-394>
- Sánchez-Muros MJ, Barroso FG, and Manzano-Agugliaro F (2014). Insect meal as renewable source of food for animal feeding: A review. *Journal of Cleaner Production*, 65: 16-27. DOI: <https://www.doi.org/10.1016/j.jclepro.2013.11.068>
- Schiavone A, Dabbou S, Petracci M, Zampiga M, Sirri F, Biasato I, and Gasco L (2019). Black soldier fly defatted meal as a dietary protein source for broiler chickens: Effects on carcass traits, breast meat quality and safety. *Animal*, 13(10): 2397-2405. DOI: <https://www.doi.org/10.1017/S1751731119000685>
- Chick feed (starter). Jakarta: National standardization agency. Available at: https://bsn.go.id/uploads/download/23kepka123_7_masa_transisi_sni_revisi_8173-2022_rev.pdf
- Sibbald IR (1980). Metabolizable energy in poultry nutrition. *BioScience*, 30: 736-741. DOI: <https://www.doi.org/10.2307/1308333>
- Sibbald IR and Wolynetz MS (1985). Relationships between estimates of bioavailable energy made with adult cockerels and chicks: Effects of feed intake and nitrogen retention. *Poultry Science*, 64(1): 127-138. DOI: <https://www.doi.org/10.3382/ps.0640127>
- Surendra KC, Tomberlin JK, van Huis A, Cammack JA, Heckmann LHL, and Khanal SK (2020). Rethinking organic wastes bioconversion: Evaluating the potential of the black soldier fly (*Hermetia illucens* (L.) (Diptera: Stratiomyidae) (BSF). *Waste Management*, 117: 58-80. DOI: <https://www.doi.org/10.1016/j.wasman.2020.07.050>
- Tan TS, Chin HY, Tsai ML, and Liu CL (2015). Structural alterations, pore generation, and deacetylation of α - and β -chitin submitted to steam explosion. *Carbohydrate Polymers*, 122: 321-328. DOI: <https://www.doi.org/10.1016/j.carbpol.2015.01.016>
- Utama CS, Cahya RI, and Sulistiyanto B (2024). Utilization of dietary maggot frass on the performance, carcass percentage, digestive organs, and economic value of Muscovy ducks. *Tropical Animal Science Journal*, 47(1): 104-111. DOI: <https://www.doi.org/10.5398/tasj.2024.47.1.104>
- Utama CS, Sulistiyanto B, and Kismiyati S (2017). The effects of water addition and steaming duration on starch composition of wheat pollard. *Reaktor*, 17(4): 221-225. Available at: <http://ejournal.undip.ac.id/index.php/reaktor/>
- Utama CS, Zuprizal, Z, Hanim C, and Wihandoyo W (2020). Effects of probiotic, prebiotic, and synbiotic mixed culture based on wheat pollard on productivity of kampung's chicken. *Indonesian Journal of Animal and Veterinary Sciences*, 25(4): 196-205. Available at: <https://medpub.appertani.org/index.php/jitv/article/download/2499/2183>
- Warisman, Agusnar H, Tafsir M, and Yunilas (2024). Effect of organic acid and steaming treatments on chitin content and nutritional quality of *Hermetia illucens* larvae as broiler feed. *Journal of Animal and Plant Sciences*, 34(6): 1606-1615. DOI: <https://www.doi.org/10.36899/JAPS.2024.6.0836>

Publisher's note: [Scienceline Publication](#) Ltd. remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025