

## Development of a Duplex Real-time PCR for Differentiation of *Salmonella* Typhimurium and Monophasic Serovars

Amany Abd El -Lattief<sup>1</sup>, Sherif Marouf<sup>2</sup>, Amany El - Bialy<sup>1</sup> and Jakeen El - Jakee<sup>2\*</sup>

<sup>1</sup>Animal Health Research Institute, Doki, Giza, Egypt

<sup>2</sup>Microbiology Department, Faculty of Veterinary Medicine, Cairo University, Egypt

\*Corresponding author's E-mail [jeljakee@yahoo.com](mailto:jeljakee@yahoo.com); ORCID: 0000-0002-5299-3783

Received: 26 Feb. 2020

Accepted: 05 Apr. 2020

### ABSTRACT

*Salmonella* Typhimurium is the most *Salmonella* serovar causing acute gastroenteritis and diarrhea. Serovar 1, 4, [5], 12: i:- is considered a monophasic variant of *S. Typhimurium* that threaten public health. Fifty-eight serologically confirmed *Salmonella* strains were investigated by PCR using *16S rRNA* and *fliC* genes. All 58 strains harbored *16S rRNA* while 21 strains harbored *fliC* gene that included *S. Typhimurium* (12), *S. Kentucky* (6), *Salmonella* variant strain serotype 1, 4, [5],12:i:- (1), *S. Lagos* (1), and *S. Kedougou* (1). A duplex TaqMan real-time PCR was performed for differentiating between biphasic *S. Typhimurium* and monophasic serovar 1, 4, [5], 12:i:- using *fljB1, 2* and *fliB/IS200* in the *fliA-fliB* intergenic region. Ten out of twelve *S. Typhimurium* harbored *fljB 1, 2*, while *Salmonella* variant strain serotype 1, 4, [5], 12:i:- lacked this gene. Thirteen strains (12 *S. Typhimurium* and the variant strain serotype 1, 4, [5], 12:i:-) were positive for *fliB/IS200* that is a specific gene for *S. Typhimurium* (biphasic and monophasic). The result of duplex TaqMan real-time PCR indicated that 10 *S. Typhimurium* strains were biphasic while two *S. Typhimurium* strains and the variant strain serotype 1, 4, [5], 12:i:- lack *fljB1,2* and had *fliB/IS200* were monophasic *S. Typhimurium*. It is noticed that prolonged subculture and repeat phase inversion method leads to the formation of flakes that in turn cause wrongly serotype identification, therefore, real-time PCR is rapid and can be used for identifying and differentiating between biphasic and monophasic *S. Typhimurium*.

**Key words:** Biphasic and monophasic *S. Typhimurium*, *flj* gene, Real-time PCR, *Salmonella*.

### INTRODUCTION

*Salmonella enterica* is zoonotic bacteria transmitted through the food chain and is an important cause of disease in humans (Osman et al., 2014a; Shaw et al., 2018). It is the second leading cause of bacterial foodborne illness (Foley et al., 2008; Persad and LeJeune, 2018). The genus *Salmonella* has a large number of serotypes that differ in pathogenicity and host specificity. Despite the widespread use of serotyping, it has deficiencies that limit its utility, including that it often takes three or more days to give a result and approximately 5-8% of isolates are partially typed.

In addition, prolonged subculture can affect the antigenic properties of strains. Highly trained laboratories are required to type strains accurately, also high costs of producing and validity specific antisera to rare antigens are problematic (Kim et al., 2006). Delay caused by identification can hinder the response to a disease outbreak and/ or epidemiological surveillance. Therefore, various studies have been explored alternative assays to

differentiate *Salmonella* isolates, such as the search for genes that can be used as potential molecular substitutes for serotyping. However, the genes tested so far have often yielded inconsistent results (Osman et al., 2014b; Hua Zou et al., 2016). Real-time PCR for detection of *Salmonella* has been brought to inter-laboratory trial, the results of which support their use as international standard methods (Malorny et al., 2007).

Two genomic sites, *16S rRNA* and *fliC* gene have been reported as candidates suitable for common and specific detection of Genus *Salmonella*, and *S. Typhimurium*, respectively by real-time PCR (Imre et al., 2005). The *16SrRNA* can be used for the rapid and multiple detections of the 16 pathogenic bacteria frequently isolated from contaminated foods that are important for food safety (Shin et al., 2016). The 16S ribosomal RNA (rRNA), approximately 1500 nucleotides in length of the prokaryotic ribosome, provides sufficient highly-conserved sequences to design the probes for developing microbial detection (Woo et al., 2003). The

*fliC* gene codes for the Hi antigen of *Salmonella* targeting the *fliC-i* allele greatly increases the specificity for *S. Typhimurium* identification (Pathmanathan et al., 2003).

*S. Typhimurium*, according to the White-Kaufmann-Le Minor serotyping scheme (Grimont and Weill, 2007), exhibits the antigenic formula 1, 4, [5], 12:i:1, 2, where “i” and “1,2” are the first and second flagellar antigens expressed by the bacterium at different times, hence the serotype description as biphasic (Soyer et al., 2009). Antigenic variants that lack either the first or second H antigens or both have been described. In recent years isolates with antigenic formula 1, 4, [5], 12:i:- have become increasingly important as a public health risk and more frequently recovered from humans and food-producing animals (Hopkins et al., 2010). The European Food Safety Authority (EFSA, 2010) recently recommended the confirmation of the serological identification of monophasic *S. 1, 4, [5], 12:i:-* strains using a polymerase chain reaction (PCR) protocol based on the detection of *fljB* gene and the *fliA-B* intergenic region. The *fljB*1,2 gene codes for second phase flagellar antigen present in *S. Typhimurium*. Indeed, all serovar *Typhimurium* strains and its monophasic/ nonmotile variants have an IS200 fragment of 1 kb in the *fliA-B* intergenic region, which is not detected in the other serovars. Within the flagellin gene cluster of *Salmonella Typhimurium* carries a conserved IS200 insertion sequence located downstream of the flagellin N-methylase gene (*fliB*) and upstream of the flagellar biosynthesis sigma factor gene (*fliA*), this element found in *Salmonella Typhimurium* and its variant (Burnens et al., 1997). Several studies have reported DNA sequences for *Salmonella* flagellin genes. As of June 2003, 74 complete or partial *Salmonella fliC* alleles and 25 complete or partial *Salmonella fljB* allele sequences had been documented in GenBank release no. 132, excluding complete genome sequences.

Thus, this study aimed, first, to confirm *Salmonella* strains using *16SrRNA* gene and *S. Typhimurium* using *fliC* by Syber green-based real-time PCR, and second, to differentiate between *S. Typhimurium* and monophasic serovar 1, 4, [5], 12:i:- using *fljB*1,2 and IS200 in the *fliA-fliB* region using TaqMan real-time PCR.

## MATERIALS AND METHODS

### Strains

A total of 58 *Salmonellae* isolates recovered from chicken in previous work (Abd El-Lattief, 2014), was identified serologically by slide agglutination test

according to White-Kauffmann le minor scheme (Grimont and Weill, 2007) using SIFIN antisera, Berlin, kindly obtained from Serology Unit, Animal Health Research Institute.

### Phase inversion method

According to ISO/TR6579 (2014), specific phase inversion antiserum was added to a swarm agar medium (SIFIN) and the *Salmonella* strain was spot inoculated on the plate. The agar medium shall be sufficiently soft for motile *Salmonella* to swarm over the medium. Slide agglutination test was performed from periphery of the plate after incubation at 37°C for 24 hrs.

### Duplex Syber green real-time PCR

For the detection of genus *Salmonella* and *S. Typhimurium*, DNA was extracted from the strains according to QIAamp DNA mini kit instructions (Soumet et al., 1999 and Yang et al., 2014). SYBR Green real-time PCR was performed using oligonucleotide primers (Table 1) and Quantitect SYBR green PCR kit containing 1ml 2xQuantiTect SYBR Green PCR Master Mix, 2ml RNase-Free Water.

**Table 1.** Oligonucleotide primers used in this study for detection of genus *Salmonella* using *16SrRNA* and *fliC* genes

| Target gene | Primer sequence (5'-3')   | Reference           |
|-------------|---------------------------|---------------------|
| 16S rRNA    | F: CAGAAGAAGCACCGGCTAACTC | Yang et al., 2014   |
|             | R: GCGCTTACGCCAGTAATT     |                     |
| <i>fliC</i> | F: CGGTGTGCCAGGTTGGTAAT   | Soumet et al., 1999 |
|             | R: ACTCTTGCTGGCGGTGCGACTT |                     |

F: forward, R: reverse

**Table 2.** Oligonucleotide primers and probes used for differentiating between biphasic *Salmonella Typhimurium* and monophasic serovar 1, 4, [5], 12:i:- using *fljB*1,2 and *fliB/IS200* in the *fliA-fliB* intergenic region.

| Target            | Primer sequence (5'-3') and probe   | Reference                |
|-------------------|---|--------------------------|
| <i>fljB</i> 1, 2  | F: TGT TAC TAT TGG TGG CTT TAC TGG  | Prendergast et al., 2013 |
|                   | R: CAG CAG GCA TTG TGG TCT TAG  |                          |
| <i>fliB/IS200</i> | FAM- CGC CAG CCG CAA GGG TTA CTG TAC – TAMRA                              | Prendergast et al., 2013 |
|                   | R: AAC GCT TGT CTT CGG TAT TTG G CY5-TCG GGT GTG CGC TAA GCT CTT TT -BHQ1 |                          |

F: forward, R: reverse

### Differentiation of *S. Typhimurium* and monophasic 1, 4, [5], 12:i-by TaqMan real-time PCR

TaqMan real-time PCR was performed according to Prendergast et al. (2013) using oligonucleotide primers and probes presented in table 2, and the Quantitect probe real-time PCR kit (Qiagen).

#### *fliC* sequencing

*fliC* was sequenced using *fliC* primer presented in table 1. A purified PCR product was sequenced in the way of the forward and/ or reverse directions on an Applied Biosystems 3130 automated DNA Sequencer (ABI, 3130, USA), using a ready reaction Bigdye Terminator V3.1 cycle sequencing kit (Perkin-Elmer/Applied Biosystems, Foster City, CA).

### Phylogenetic analysis

A comparative analysis of sequences was performed using the CLUSTAL W multiple sequence alignment program, version 1.83 of MegAlign module of Lasergene DNA Star software Pairwise, which was designed by Thompson et al. (1994) and phylogenetic analyses were done using maximum likelihood, neighbor-joining and maximum parsimony in MEGA6 (Tamura et al., 2013).

## RESULTS

### Serotyping of *Salmonella*

The Serotyping of the 58 *Salmonella* strains was confirmed and the result is presented in table 3.

**Table 3.** Antigenic structure of all *Salmonella* strains recovered using slide agglutination test.

| No  | Name                  | Serotyping                    | No  | Name                   | Serotyping                                      |
|-----|-----------------------|-------------------------------|-----|------------------------|---|
| 1-  | <i>S. Kentucky</i>    | 8,20 :i:z <sub>6</sub>        | 30- | <i>S. Washington</i>   | 13,22 :m,t:-                                    |
| 2-  | <i>S. Lagos</i>       | 1,4,5,12:i:1,5                | 31- | <i>S. Newport</i>      | 6,8,20 :e,h :1,2                                |
| 3-  | <i>S. Typhimurium</i> | 1,4,[5],12:i:1,2              | 32- | <i>S. Enteritidis</i>  | 1,9,12 : g,m:-                                  |
| 4-  | <i>S. Typhimurium</i> | 1,4,[5],12:i:1,2              | 33- | <i>S. Rissen</i>       | 6,7,14 :f,g:-                                   |
| 5-  | <i>S. Taksony</i>     | 1,3,19: i: z <sub>6</sub>     | 34- | <i>S. Labadi</i>       | 8,20 :d: z <sub>6</sub>                         |
| 6-  | <i>S. Derby</i>       | 1,4,[5],12:f,g:-              | 35- | <i>S. Enteritidis</i>  | 1,9,12 : g,m: -                                 |
| 7-  | <i>S. Rissen</i>      | 6,7,14:f,g:-                  | 36- | <i>S. Senftenberg</i>  | 1,3,19: g,[s],t :-                              |
| 8-  | <i>S. Typhimurium</i> | 1,4,[5],12:i:1,2              | 37- | <i>S. Cerro</i>        | 6,14,18 :z <sub>4</sub> ,z <sub>23</sub> :[1,5] |
| 9-  | <i>S. Anatum</i>      | 3,{10} {15} {15,34} :e,h: 1,6 | 38- | <i>S. Virginia</i>     | 8: d :1,2                                       |
| 10- | <i>S. Typhimurium</i> | 1,4,[5],12:i:1,2              | 39- | <i>S. Papuana</i>      | 6,7 :r :e,n,z <sub>15</sub>                     |
| 11- | <i>S. Paratyphi A</i> | 1,2,12: a: [1,5]              | 40- | <i>S. Typhimurium</i>  | 1,4,[5],12:i:1,2                                |
| 12- | <i>S. Paratyphi B</i> | 1,4,[5],12 :b: 1,2            | 41- | <i>S. Typhimurium</i>  | 1,4,[5],12:i:1,2                                |
| 13- | <i>S. Kedougou</i>    | 1,13,23: i :l,w               | 42- | <i>S. Kentucky</i>     | 8,20: i: z <sub>6</sub>                         |
| 14- | <i>S. Labadi</i>      | 8,20 :d: z <sub>6</sub>       | 43- | <i>S. Typhimurium</i>  | 1,4,[5],12:i:1,2                                |
| 15- | <i>S. Poona</i>       | 1,13,22: z: 1,6               | 44- | <i>S. Enteritidis</i>  | 1,9,12 :g,m: -                                  |
| 16- | <i>S. Typhimurium</i> | 1,4,[5],12:i:1,2              | 45- | <i>S. Virginia</i>     | 8:d:1,2   |
| 17- | <i>S. Kentucky</i>    | 8,20: i :z <sub>6</sub>       | 46- | <i>S. Kentucky</i>     | 8,20: i :z <sub>6</sub>                         |
| 18- | <i>S. Anatum</i>      | 3,{10}{15}{15,34} :e,h: 1,6   | 47- | <i>S. Washington</i>   | 13,22 :m,t:-                                    |
| 19- | <i>S. Goldcoast</i>   | 6,8 :r :l,w                   | 48- | <i>S. Enteritidis</i>  | 1,9,12:g,m: -                                   |
| 20- | <i>S. Enteritidis</i> | 1,9,12:g,m:-                  | 49- | <i>S. Newlands</i>     | 3,{10} {15 ,34}:e,he,n,x:-                      |
| 21- | <i>S. Infantis</i>    | 6,7,14: r :1,5                | 50- | <i>S. Gallinarum</i>   | 1,9,12 :-: -                                    |
| 22- | <i>S. Gallinarum</i>  | 1,9,12: - :-                  | 51  | <i>S. Agama</i>        | 4,12:i:1,6                                      |
| 23- | <i>S. Gallinarum</i>  | 1,9,12: - :-                  | 52- | <i>S. Kentucky</i>     | 8,20: i: z <sub>6</sub>                         |
| 24- | <i>S. Hadar</i>       | 6,8 :z <sub>10</sub> :e,n,x   | 53- | <i>S. Kentucky</i>     | 8,20: i :z <sub>6</sub>                         |
| 25- | <i>S. Virchow</i>     | 6,7,14: r :1,2                | 54- | <i>S. Typhimurium</i>  | 1,4,[5],12:i:1,2                                |
| 26- | <i>S. Virchow</i>     | 6,7,14: r :1,2                | 55- | <i>S. Typhimurium</i>  | 1,4,[5],12 :i :1111,2                           |
| 27- | <i>S. Hadar</i>       | 6,8 :z <sub>10</sub> :e,n,x   | 56- | <i>S. Typhimurium</i>  | 1,4,[5],12:i:1,2                                |
| 28- | <i>S. Bardo</i>       | 8 :e,h:1,2                    | 57- | Partial identification | 1,4,[5],12:i:-                                  |
| 29- | <i>S. Montevideo</i>  | 6,7,14 :g,m,s :-              | 58- | <i>S. Typhimurium</i>  | 1,4,[5],12:i:1,2                                |

**Syber Green real-time duplex PCR**

All 58 strains belonged to genus *Salmonella* were positive by SYBER green real-time PCR using *16S rRNA*. The specificity of the reaction was confirmed by melting temperature (Tm) which was consistently specific for amplicon obtained; the mean peaks Tm obtained. The negative control did not show peaks in the Tm when subjected to 40 cycles of amplification. Twenty-one *Salmonella* strains harbored *fliC* gene, including *S. Typhimurium* (12), *S. Kentucky* (6), *S. Lagos* (1), *S. Kedougou* (1) and partial identified strain *S* 1,4, [5],12:i:- which possess first flagellar i antigen (Table 4; figures 1 and 2). A total of 15 strains were positive for *fljB* 1,2. Ten strains of *Salmonella* Typhimurium and serovars Paratyphi A (1) & Paratyphi B (1) & Newport (1) and Virginia (2) harbored *fljB*1,2, while strain no.57 with antigenic formula 1, 4, [5], 12:i:- and two strains *Salmonella* Typhimurium

lacked this gene. Concerning *fli B/IS200*, the 13 strains possess *fliB/IS200* (12 *S. Typhimurium* and the variant strain *S* 1, 4, [5], 12:i:-) (Table 5 and figure 3).

***fli C* Sequencing**

Individual *Salmonella* serotypes usually alternate between the production of 2 antigenic forms of flagella, termed phase 1 and phase 2, each specified by separate structural genes, *fliC* and *fljB* 1, 2. Sequencing of *fliC* gene based on the nucleotide sequence of *S. Typhimurium*13311 referenced in GenBank illustrated that the biphasic *S. Typhimurium* strain was recorded in GenBank as *S. Typhimurium* Egy 1 with accession number Mk103394 and the monophasic strain as *S. Typhimurium* Egy 2 with accession number MK 103395. The amino acid sequence of the *fliC* gene in the two isolates showing greater than 98% identity.


**Table 4.** Detection of *16SrRNA* and *fliC* genes in *Salmonella* serovars using duplex Syber green real-time PCR

| No. | Name                  | 16S RNA | fliC | No. | Name                  | 16S rRNA | fliC |
|-----|-----------------------|---------|------|-----|-----------------------|----------|------|
| 1   | <i>S. Kentucky</i>    | +       | +    | 30  | <i>S. Washington</i>  | +        | -    |
| 2   | <i>S. Lagos</i>       | +       | +    | 31  | <i>S. Newport</i>     | +        | -    |
| 3   | <i>S. Typhimurium</i> | +       | +    | 32  | <i>S. Enteritidis</i> | +        | -    |
| 4   | <i>S. Typhimurium</i> | +       | +    | 33  | <i>S. Rissen</i>      | +        | -    |
| 5   | <i>S. Taksony</i>     | +       | -    | 34  | <i>S. Labadi</i>      | +        | -    |
| 6   | <i>S. Derby</i>       | +       | -    | 35  | <i>S. Enteritidis</i> | +        | -    |
| 7   | <i>S. Rissen</i>      | +       | -    | 36  | <i>S. Senftenberg</i> | +        | -    |
| 8   | <i>S. Typhimurium</i> | +       | +    | 37  | <i>S. Cerro</i>       | +        | -    |
| 9   | <i>S. Anatum</i>      | +       | -    | 38  | <i>S. Virginia</i>    | +        | -    |
| 10  | <i>S. Typhimurium</i> | +       | +    | 39  | <i>S. Papuana</i>     | +        | -    |
| 11  | <i>S. Paratyphi A</i> | +       | -    | 40  | <i>S. Typhimurium</i> | +        | +    |
| 12  | <i>S. Paratyphi B</i> | +       | -    | 41  | <i>S. Typhimurium</i> | +        | +    |
| 13  | <i>S. Kedougou</i>    | +       | +    | 42  | <i>S. Kentucky</i>    | +        | +    |
| 14  | <i>S. Labadi</i>      | +       | -    | 43  | <i>S. Typhimurium</i> | +        | +    |
| 15  | <i>S. Poona</i>       | +       | -    | 44  | <i>S. Enteritidis</i> | +        | -    |
| 16  | <i>S. Typhimurium</i> | +       | +    | 45  | <i>S. Virginia</i>    | +        | -    |
| 17  | <i>S. Kentucky</i>    | +       | +    | 46  | <i>S. Kentucky</i>    | +        | +    |
| 18  | <i>S. Anatum</i>      | +       | -    | 47  | <i>S. Washington</i>  | +        | -    |
| 19  | <i>S. Goldcoast</i>   | +       | -    | 48  | <i>S. Enteritidis</i> | +        | -    |
| 20  | <i>S. Enteritidis</i> | +       | -    | 49  | <i>S. Newlands</i>    | +        | -    |
| 21  | <i>S. Infantis</i>    | +       | -    | 50  | <i>S. Gallinarum</i>  | +        | -    |
| 22  | <i>S. Gallinarum</i>  | +       | -    | 51  | <i>S. Agama</i>       | +        | -    |
| 23  | <i>S. Gallinarum</i>  | +       | -    | 52  | <i>S. Kentucky</i>    | +        | +    |
| 24  | <i>S. Hadar</i>       | +       | -    | 53  | <i>S. Kentucky</i>    | +        | +    |
| 25  | <i>S. Virchow</i>     | +       | -    | 54  | <i>S. Typhimurium</i> | +        | +    |
| 26  | <i>S. Virchow</i>     | +       | -    | 55  | <i>S. Typhimurium</i> | +        | +    |
| 27  | <i>S. Hadar</i>       | +       | -    | 56  | <i>S. Typhimurium</i> | +        | +    |
| 28  | <i>S. Bardo</i>       | +       | -    | 57  | 1,4,[5],12:i:-        | +        | +    |
| 29  | <i>S. Montevideo</i>  | +       | -    | 58  | <i>S. Typhimurium</i> | +        | +    |


**Table 5.** Detection of *fljB1,2* and *fliB/IS200* in *Salmonella* serovars using duplex TaqMan real-time PCR

| No. | Name                  | <i>fljB1,2</i> | <i>fliB/IS200</i> | No. | Name                  | <i>fljB1,2</i> | <i>fliB /IS200</i> |
|-----|-----------------------|----------------|-------------------|-----|-----------------------|----------------|--------------------|
| 1   | <i>S. Kentucky</i>    | -              | ND                | 30  | <i>S. Washington</i>  | -              | ND                 |
| 2   | <i>S. Lagos</i>       | -              | ND                | 31  | <i>S. Newport</i>     | +              | ND                 |
| 3   | <i>S. Typhimurium</i> | +              | +                 | 32  | <i>S. Enteritidis</i> | -              | ND                 |
| 4   | <i>S. Typhimurium</i> | +              | +                 | 33  | <i>S. Rissen</i>      | -              | ND                 |
| 5   | <i>S. Taksony</i>     | -              | ND                | 34  | <i>S. Labadi</i>      | -              | ND                 |
| 6   | <i>S. Derby</i>       | -              | ND                | 35  | <i>S. Enteritidis</i> | -              | ND                 |
| 7   | <i>S. Rissen</i>      | -              | ND                | 36  | <i>S. Senftenberg</i> | -              | ND                 |
| 8   | <i>S. Typhimurium</i> | +              | +                 | 37  | <i>S. Cerro</i>       | -              | ND                 |
| 9   | <i>S. Anatum</i>      | -              | ND                | 38  | <i>S. Virginia</i>    | +              | ND                 |
| 10  | <i>S. Typhimurium</i> | +              | +                 | 39  | <i>S. Papuana</i>     | -              | ND                 |
| 11  | <i>S. Paratyphi A</i> | +              | ND                | 40  | <i>S. Typhimurium</i> | +              | +                  |
| 12  | <i>S. Paratyphi B</i> | +              | ND                | 41  | <i>S. Typhimurium</i> | +              | +                  |
| 13  | <i>S. Kedougou</i>    | -              | ND                | 42  | <i>S. Kentucky</i>    | -              | ND                 |
| 14  | <i>S. Labadi</i>      | -              | ND                | 43  | <i>S. Typhimurium</i> | +              | +                  |
| 15  | <i>S. Poona</i>       | -              | ND                | 44  | <i>S. Enteritidis</i> | -              | ND                 |
| 16  | <i>S. Typhimurium</i> | +              | +                 | 45  | <i>S. Virginia</i>    | +              | ND                 |
| 17  | <i>S. Kentucky</i>    | -              | ND                | 46  | <i>S. Kentucky</i>    | -              | ND                 |
| 18  | <i>S. Anatum</i>      | -              | ND                | 47  | <i>S. Washington</i>  | -              | ND                 |
| 19  | <i>S. Goldcoast</i>   | -              | ND                | 48  | <i>S. Enteritidis</i> | -              | ND                 |
| 20  | <i>S. Enteritidis</i> | -              | ND                | 49  | <i>S. Newlands</i>    | -              | ND                 |
| 21  | <i>S. Infantis</i>    | -              | ND                | 50  | <i>S. Gallinarum</i>  | -              | ND                 |
| 22  | <i>S. Gallinarum</i>  | -              | ND                | 51  | <i>S. Agama</i>       | -              | ND                 |
| 23  | <i>S. Gallinarum</i>  | -              | ND                | 52  | <i>S. Kentucky</i>    | -              | ND                 |
| 24  | <i>S. Hadar</i>       | -              | ND                | 53  | <i>S. Kentucky</i>    | -              | ND                 |
| 25  | <i>S. Virchow</i>     | -              | ND                | 54  | <i>S. Typhimurium</i> | +              | +                  |
| 26  | <i>S. Virchow</i>     | -              | ND                | 55  | <i>S. Typhimurium</i> | -              | +                  |
| 27  | <i>S. Hadar</i>       | -              | ND                | 56  | <i>S. Typhimurium</i> | -              | +                  |
| 28  | <i>S. Bardo</i>       | -              | ND                | 57  | <i>S. Typhimurium</i> | -              | +                  |
| 29  | <i>S. Montevideo</i>  | -              | ND                | 58  | <i>S. Typhimurium</i> | +              | +                  |


ND: Not detected




**Figure 1.** Syber green real-time PCR targeting *16S rRNA* gene for 58 *Salmonella* strains isolated from chickens (fluorescence chart and melting curve). A) Fluorescence chart for strains number 1 to 29. B) Fluorescence chart for strains number 30 to 58. (Amplification plots represent the accumulation of product over the duration of real-time PCR). C) Melting curve for strain number 1 to 29. D) Melting curve for strain number 30 to 58. Melting curve provides representation of the PCR product after the amplification process, A single peak indicates a positive sample. All 58 strains isolated from chickens were positive for *16S rRNA* gene. The specificity of the reaction was confirmed by the melting temperature. The mean peak temperature obtained was 80.55 °C.



**Figure 2.** Syber green real-time PCR targeting *fliC* gene for 58 *Salmonella* strains isolated from chicken (fluorescence chart and melting curve). A) Fluorescence chart for strains number 1 to 29. B) Fluorescence chart for strains number 30 to 58 (Amplification plots represent the accumulation of product over the duration of real-time PCR). C) Melting curve for strain number 1 to 29, where strains number 1, 2, 3, 4, 8, 10, 13, 16 and 17 gave positive results. D) Melting curve for strain number 30 to 58, where strains number 40, 41, 42, 43, 46, 52, 53, 54, 55, 56, 57 and 58 gave positive result. (Melting curve provide representation of the PCR product after the amplification process. A single peak indicates a positive sample. Twenty-one *Salmonella* strains harbored *fliC* gene. The specificity of the reaction was confirmed by melting temperature, the mean peak temperature obtained was 85.65 °C.



**Figure 3.** TaqMan real-time PCR amplification chart for *fliB*1,2 gene among 58 *Salmonella* strains isolated from chickens. Typical amplification curves given for positive samples. Fifteen strains (number 3, 4, 8, 10, 11, 12, 16, 31, 38, 40, 41, 43, 45, 54 and 58) gave positive results.



**Figure 4.** TaqMan real-time PCR amplification chart for *fliB/IS200* gene among 13 *Salmonella* isolates (12 *S. Typhimurium* and the variant strain serotype 1, 4, [5], 12:i:-) (*fliB/IS200* is a specific gene for *S. Typhimurium* biphasic and monophasic). Thirteen strains (number 3, 4, 8, 10, 16, 40, 41, 43, 54, 55, 56, 57 and 58) gave positive results.

## DISCUSSION

Serological identification of 58 *Salmonella* strains was confirmed by slide agglutination test and the antigenic structure is demonstrated in table 3. Failure to identify the complete antigenic formula prevents the unequivocal identification of serovars even after phase inversion

method. The strain was considered monophasic when phase inversion method was repeated at least three times without getting expression of phase 2 flagellar antigen as shown in strain number 57 with antigenic formula S.1, 4, [5], 12:i:-. Grimont and Weil (2007) mentioned that S.1, 4, [5], 12:i:- does not appear in the White-Kaufmann-Le Minor scheme and appears to be a monophasic variant of other biphasic serovars, which have lost phase 2 flagellin or the necessary switching mechanism of phase variation. Seven serovars of *S. enterica* subsp. *enterica* with the same O and phase 1 H antigens are possible ancestors of this serovar, including *S. Typhimurium*, *S. Lagos*, *S. Agama*, *S. Farsta*, *S. Tsevie*, *S. Gloucester*, and *S. Tumodi*. Among these, *S. Typhimurium* monophasic S.1, 4, [5], 12:i:- is commonly isolated from humans, animals, and the environment.

In recent years, many studies try to establish methods that can reduce the time for the detection and identification of salmonellae. Detection of bacteria by conventional methods is time-consuming and allows the detection of viable one only (Kim et al., 2006).

The use of PCR has emerged as an approach to overcome these problems. The exploration of gene targets for evaluation of absence and presence of bacteria is still a matter of importance. Several genes *invA*, *fimA*, and *aceK* were used for identification of genus *Salmonella* (O'Regan et al., 2008). The duplex Syber green real-time PCR was applied for detection of genus *Salmonella* and the most common serovar *S. Typhimurium* based on melting Temp (TM) and Curve analysis using *16S rRNA* and *fliC* genes respectively. *16S rRNA* not only allow the presence of bacteria to be proved but also would give information on gene expression. However, the expression of rRNA is tightly depend on physiological status of bacteria (Imre et al., 2005). In this study all 58 *Salmonella* strains harbor *16S rRNA* (Table 4 and figure 1).

*16SrRNA* gene sequences contain hypervariable regions that can allow species-specific signature sequences important for identification of bacteria. The *16SrRNA* gene is used as the standard for classification and identification of bacteria because it is present in most microbes and shows proper changes. *16SrRNA* gene sequences for most bacteria are available on public databases such as NCBI (Pereira, 2010). Attractive potential uses of *16SrRNA* gene sequence informatics for providing genus and species identification.

The *fliC* target is specific for the phase-1 flagellar antigen i that encoded by serovars Typhimurium. In the present study twenty one strains possess *fliC* gene serovars Typhimurium (12), Kentucky (6), Kedougou (1), Lagos



(1) and *S.* 1,4, [5], 12:i:-(1) (Table 4 and figure 2). O' Regan *et al.* (2008) reported that the i antigen is also expressed in uncommon serotypes such as Aberdeen, Bergen, and Kedougou. The structural flagellin gene *fliC* was present in all isolates of serovars Typhimurium and Kentucky (full length) and in all isolates of serovars Heidelberg, Hadar, and Enteritidis (partial length) (Dhanani *et al.*, 2015).

Most *S. enterica* serovar Typhimurium possess two different flagellin proteins, including FliC (phase 1) and FljB (phase 2), which are encoded by the genes *fliC* and *fljB*, respectively. European Food Safety Authority (EFSA) (2010) applied a conventional PCR protocol to confirm the absence of 2<sup>nd</sup> phase antigen. A real-time PCR assay was used to differentiate *S.* Typhimurium monophasic variants from biphasic *S.* Typhimurium and from other variants (Anon, 2010; Tennant *et al.*, 2010).

Fifteen isolates are positive for *fljB1,2* *S.* Typhimurium (10), *S.* Paratyphi A(1), *S.* Paratyphi B(1), *S.* Newport (1) and *S.* Virginia (2) (Table 5 and figure 3). This result agree with that published by Bugarel *et al.* (2012) who reported that the second gene codes for the phase 2 flagellar antigen *fljB1,2* is present in *S.* Typhimurium and other serovars such as *S.* Coeln, *S.* Haifa, *S.* Heidelberg, *S.* Paratyphi B, *S.* Saintpaul and *S.* Stanley. This marker is absent in monophasic *S.* Typhimurium. Two serologically identified *S.* Typhimurium strains no. 55, 56 don't possess *fljB1,2* that could be explained by repeat phase inverted method leads to formation of flakes which may lead to misidentification or wrongly identified strains.

Flagellar phase variation is formed by inversion of the genetic region called the H segment, which have the *hin* gene encoding for DNA invertase and the promoter for the *fljB* gene. The *fljB* constitutes an operon with the *fljA* gene, which encodes a negative regulator of *fliC* expression. FljA binds to the operator region of FliC mRNA and inhibits its translation, leading to the rapid degradation of FliC mRNA. When the H segment is in the "on" state, both *fljB* and *fljA* are transcribed, lead to synthesis of phase 2 flagellin and inhibition of phase 1 flagellin. However, when the H segment is switched to the "off" state, neither *fljB* nor *fljA* are transcribed, resulting in the synthesis of phase 1 flagellin only (Ido *et al.*, 2014).

The location of IS200 between the genes *fliA* and *fliB* can be used as a specific marker for *S.* Typhimurium. The amplicon sizes from the *fliA*-*fliB* intergenic regions from *S.* Typhimurium and other serovars were expected to be 1000 and 250 bp, respectively. TaqMan real-time PCR could successfully detected *S.* 1, 4, [5], 12:i:- isolates that

yield 1000-bp amplicon with conventional PCR. These data suggest that *S.* 1, 4, [5], 12:i:- is a monophasic variant of *S.* Typhimurium (Burnens *et al.*, 1997). Also, they reported that within the flagellin gene cluster of *Salmonella*, *S.* Typhimurium carries a conserved IS200 insertion sequence located downstream of the flagellin N-methylase gene (*fliB*) and upstream of the flagellar biosynthesis sigma factor gene (*fliA*). In the present study ten strains yield positive result with *fliC*, *fljB1,2* and *fliB*/IS200 were biphasic *Salmonella* Typhimurium meanwhile 3 strains harbored the *fliC* and *fliB*/IS200 were monophasic strains *S.* 1, 4, [5], 12:i:- (Table 6).

During recent years the cost of sequencing has been reduced dramatically making sequencing based typing more attractive. Some studies have reported DNA sequence for flagellin gene (Silverman, 1979; Joys, 1985 and De Vries, 1998). As in 2016, *fliC* sequence (partial coding sequence) has reported in GenBank with accession no DQ095491. This study reported sequencing of *fliC* gene for two strains *S.* Typhimurium and monophasic variant *S.* 1, 4, [5], 12:i:- with accession no (Mk103394) and (Mk103395), respectively.


Protein sequence is the practical process of determining the amino acid sequence of all or part of protein or peptide. About 500 naturally occurring amino acids are known, 20 only appear in the genetic code there are termed as codons are always 3 Base pairs (nucleotides). In this study, amino acid sequence were applied for the *fliC* gene. In the location 14-19 sequence TNGKVT was found, which is similar to sequences coded in GenBank with accession no. CP024619, LT795114, CP014979, but in other sequences reported in GenBank with accession no. CP026700, CP021462, CP028199 glutamic acid was found between GK with amino acid sequence TNGEKVT (Figure 5).


In this study, the amino acid threonine was absent at position 24 in *S.* Typhimurium Egy1 and *S.* Typhimurium Egy2, which is similar to sequences recorded in GenBank with accession no. CP014979, CP014967. While the result disagreed with sequences coded in GenBank with accession number CP007581 and DQ09549 which have threonine at position 24 between glycine and alanine.

At position 60-65 found amino acid sequence AGVTGT in *S.* Typhimurium Egy1 and *S.* Typhimurium Egy2, but in sequence coded in GenBank with accession no. LN999997 amino acid alanine at position 65 between glycine and threonine was found. Alignments show highly degree of identity. There are greater than 98% amino acid sequence identity (Figures 6 and 7). This is according to Sandjong *et al.* (2007).

**Table 6.** Comparison between results of conventional serotyping and real-time PCR for *Salmonella* Typhimurium (biphasic and monophasic strains)

| No. of isolate | Name of isolate                            | Conventional serotyping |                   |                   | Real -time PCR |                |                   |
|----------------|--|-------------------------|-------------------|-------------------|----------------|----------------|-------------------|
|                |  | O antigen               | Phase 1 H antigen | Phase 2 H antigen | <i>fliC</i>    | <i>fliB1,2</i> | <i>fliB/IS200</i> |
| 10 strains     | <i>Salmonella</i> Typhimurium (diphasic)   | 4,[5],12                | I                 | 1,2               | +              | +              | +                 |
| 3 strains      | <i>Salmonella</i> Typhimurium (monophasic) | 4,[5],12                | I                 | Not detected      | +              | -              | +                 |
| 1 strain       | Non- <i>Salmonella</i> Typhimurium         | 4,[5],12                | I                 |                   | +              | -              | -                 |
| 5strains       | Non <i>Salmonella</i> Typhimurium          | 4,[5],12                | -                 | 1,2               | -              | +              | -                 |






**Figure 5.** Amino acid sequence alignment report for *fljC* gene of two Egyptian *Salmonella* strains recorded in GenBank with accession number Mk103394 and Mk103395 for *S. Typhimurium* Egy 1 (biphasic) and *S. Typhimurium* Egy 2 (monophasic), respectively. The sequence alignment of two Egyptian strains is 100% similar to nine strains recorded in GenBank (*S. Typhimurium* BL10, *S. Typhimurium* VNB151, *S. Typhimurium* CDC H2662, *S. Typhimurium* 81741, *S. Typhimurium* USDA-ARS-USMA, *S. Typhimurium* CDC2011K-1702, *S. Typhimurium* RM9437, *S. Typhimurium* 33676 and *S. Typhimurium* SGSC2193). In the location 14-19, sequence TNGKVT was found for two Egyptian strains that matched sequences of some strains coded in GenBank with accession no. CP024619, LT795114, and CP014979, but in other strains reported in GenBank with accession no. CP026700, CP021462, and CP028199 glutamic acid was found between GK and amino acid sequence was TNGEKVT. The amino acid threonine was absent at position 24 in *S. Typhimurium* Egy1 and *S. Typhimurium* Egy2, but strains recorded in GenBank with accession no. CP007581 and DQ09549 have threonine at position 24 between glycine and alanine. At position 60-65 amino acid sequence AGVTGT was found in *S. Typhimurium* Egy1 and *S. Typhimurium* Egy2, but in a sequence coded in GenBank with accession no. LN999997 amino acid alanine was found at position 65 between glycine and threonine.

|            |    | Percent Identity |      |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |      |      |    |   |
|------------|----|------------------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|------|----|---|
|            |    | 1                | 2    | 3     | 4     | 5     | 6     | 7     | 8     | 9     | 10    | 11    | 12    | 13    | 14    | 15    | 16    | 17    | 18    | 19    | 20    | 21    | 22    | 23    | 24    | 25    | 26    | 27   |      |    |   |
| Divergence | 1  | ■                | 99.8 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.7  | 99.7 | 99.7 | 1  | CP009102.1 S. Typhimurium strain 13311  |
|            | 2  | 0.2              | ■    | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 2  | S. Typhimurium Egy 1                    |
|            | 3  | 0.2              | 0.0  | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 3  | S. Typhimurium Egy 2                    |
|            | 4  | 0.2              | 0.0  | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 4  | CP024619.1 S. Typhimurium BL10          |
|            | 5  | 0.2              | 0.0  | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 5  | LT795114.1 S. Typhimurium VNB151        |
|            | 6  | 0.2              | 0.0  | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 6  | CP014979.2 S. Typhimurium CDC H2662     |
|            | 7  | 0.2              | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 7  | CP019442.1 S. Typhimurium 81741         |
|            | 8  | 0.2              | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 8  | CP014969.1 S. Typhimurium USDA-ARS-USMA |
|            | 9  | 0.2              | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 9  | CP014967.1 S. Typhimurium CDC2011K-1702 |
|            | 10 | 0.2              | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 10 | CP012985.1 S. Typhimurium RM9437        |
|            | 11 | 0.2              | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 11 | CP012681.1 S. Typhimurium 33676         |
|            | 12 | 0.2              | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8  | 99.8 | 99.8 | 12 | AY649718.1 S. Typhimurium SGSC2193      |
|            | 13 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.7 | 99.7 | 13 | CP021462.1 S. Typhimurium UGA14e        |
|            | 14 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.7 | 99.7 | 14 | CP028199.1 S. Typhimurium CFSAN018746   |
|            | 15 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.7 | 99.7 | 15 | CP026700.1 S. Typhimurium AR_0031       |
|            | 16 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.7 | 99.7 | 16 | LT855376.1 S. Typhimurium STMU2UK       |
|            | 17 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.7 | 99.7 | 17 | CP013720.1 S. Typhimurium RM10607       |
|            | 18 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.7 | 99.7 | 18 | CP002487.1 S. Typhimurium ST4/74        |
|            | 19 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.7 | 99.7 | 19 | HM920247.1 S. Typhimurium IVRI          |
|            | 20 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 99.7 | 99.7 | 20 | AP011957.1 S. Typhimurium T000240       |
|            | 21 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 100.0 | 99.7  | 99.7 | 99.7 | 21 | FQ312003.1 S. Typhimurium SL1344        |
|            | 22 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 100.0 | 99.7  | 99.7 | 99.7 | 22 | CP001363.1 S. Typhimurium 14028S        |
|            | 23 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 100.0 | 99.7  | 99.7 | 99.7 | 23 | EF599295.1 S. Typhimurium 95            |
|            | 24 | 0.0              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | ■     | 99.7 | 99.7 | 24 | DQ095487.1 S. Typhimurium LT 2          |
|            | 25 | 0.3              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3  | 0.3  | 25 | LN999997.1 S. Typhimurium SO4698-09 I   |
|            | 26 | 0.3              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3  | 0.3  | 26 | CP007581.1 S. Typhimurium 138736        |
|            | 27 | 0.3              | 0.2  | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3   | 0.3  | 0.0  | 27 | DQ095491.1 S. Typhimurium S05917 03     |

**Figure 6.** Amino acid sequence distance performed using the CLUSTAL W multiple sequence alignment program and version 1.83 of MegAlign module of Lasergene DNASTar software Pairwise for *fliC* gene among two Egyptian *Salmonella* strains (*S. Typhimurium* Egy 1 (biphasic) and *S. Typhimurium* Egy 2 (monophasic)).



**Figure 7.** Phylogenetic analysis of *Salmonella* Typhimurium using *fliC* gene sequence performed by maximum likelihood, neighbor-joining and maximum parsimony implemented in MEGA6. The amino acid sequence of two Egyptian strain (Mk103394 and Mk103395) were closely related to sequences recorded in GenBank (CP024619 for *S. Typhimurium* BL10, LT795114 for *S. Typhimurium* VNB151, CP014979 for *S. Typhimurium* CDC H2662, CP019442 for *S. Typhimurium* 81741, CP014969 for *S. Typhimurium* USDA-ARS-USMA, CP014967 for *S. Typhimurium* CDC2011K-1702, CP012985 for *S. Typhimurium*, CP012681 for *S. Typhimurium* 33676RM9437 and AY649718 for *S. Typhimurium* SGSC2193).

## CONCLUSION

The duplex real-time PCR is a rapid and robust method for detection of genus *Salmonella* and can be used for identification and differentiation of *S. Typhimurium* and the most common variant S.1, 4, [5], 12:i:-.

## DECLARATIONS

### Acknowledgments

This study was supported by the research group of Animal Health Research Institute, Egypt

### Competing interests

The authors have declared that no competing interest exists.

### Authors' contribution

All authors contributed equally to this work

## REFERENCES

- Abd El-Lattief A (2014). Detection of multidrug resistant *Salmonellae*. master thesis, Microbiology department, Faculty of veterinary medicine, Cairo University.
- Anon (2010). Scientific opinion on monitoring and assessment of the public health risk of *Salmonella* Typhimurium-like strains. EFSA Journal, 8 (10):1826. DOI: <https://doi.org/10.2903/j.efsa.2010.1826>.
- Bugarel M, Granier SA, Bonin E, Vignaud ML, Roussel S, and Fach P (2012). Genetic diversity in monophasic (1,4,[5],12:i:- and 1,4,[5],12:i:-:1.2) and in non-motile (1,4,[5],12:i:-) variants of *Salmonella enterica* Typhimurium. Food Research International, 45 (10): 16-1024. DOI: <https://doi.org/10.1016/j.foodres.2011.06.057>
- Burnens AP, Stanley J, Sack R, Hunziker P, Brodard I and Nicolet J (1997). The flagellin N methylase gene *fliB* and an adjacent serovar-specific IS200 element in *Salmonella* Typhimurium. Microbiology, 143:1539–1547. DOI:<https://doi.org/10.1099/00221287-143-5-1539>
- De Vries N, Zwaagstra KA, Huis JHJ, Veld INT, Van Knapen F, VanZijderveld FG and Kusters JG (1998). Production of monoclonal antibodies specific for the i and 1, 2 flagellar antigens of *Salmonella Typhimurium* and characterization of their respective epitopes. Journal of Applied and Environmental Microbiology, 64(12):5033-5038. Available at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC90964/>
- Dhanani AS, Block G, Forgetta V, Topp E, Beiko R and Diarra M (2015). Genomic comparison of non-typhoidal *Salmonella enterica* serovars Typhimurium, Enteritidis, Heidelberg, Hadar and Kentucky isolates from broiler chickens. PLOS One, 10(9): 0137697e. DOI:<https://doi.org/10.1371/journal.pone.0137697>
- European Food Safety Authority (EFSA) (2010). Scientific opinion on monitoring and assessment of the public health risk of “*Salmonella* Typhimurium-like” strains. EFSA Journal, 8: 1826. DOI: <https://doi.org/10.2903/j.efsa.2010.1826>
- Foley SL, Lynne AM and Nayak R (2008). *Salmonella* challenges: Prevalence in swine and poultry and potential pathogenicity of such isolates. Journal of Animals Science, 86:149–162. DOI: <https://doi.org/10.2527/jas.2007-0464>
- Grimont PAD and Weill FX (2007). Antigenic Formulas of the *Salmonella* Serovars, 9<sup>th</sup> edition. WHO Collaborating Centre for Reference and Research on *Salmonella*. Institute Pasteur, Paris.
- Hopkins KL, Kirchner M, Guerra B, Granier SA, Lucarelli C and Porrero MC (2010). Multi resistant *Salmonella enterica* serovar 4, [5], 12:i:- in Europe: a new pandemic strain. Eurosurveillance Journal, 15: 19580. DOI: <https://doi.org/10.2807/ese.15.22.19580-en>
- Hua Zou Q, Qing Li R, Rong Liu G and Lin Liu S (2016). Genotyping of *Salmonella* with lineage-specific genes: correlation with serotyping International, Journal of Infectious Diseases, 49:134-140. DOI:<https://doi.org/10.1016/j.ijid.2016.05.029>.
- Ido N, Lee KI, Iwabuchi K, Izumiya H and Uchida I (2014). Characteristics of *Salmonella enterica* Serovar 4, [5], 12:i:- as a Monophasic Variant of Serovar Typhimurium. PLoS ONE, 9(8): e104380. DOI: <https://doi.org/10.1371/journal.pone.0104380>.
- Imre A, Olasz F and Nagy B (2005). Development of a PCR system for the characterisation of *Salmonella* flagellin genes. Acta Veterinaria Hungarica, 53:163–172. DOI: <https://doi.org/10.1556/AVet.53.2005.2.2>.
- ISO/TR6579 -3 (2014). Microbiology of food chain horizontal method for the detection, enumeration and serotyping of *Salmonella* –part3 guidelines for serotyping of *Salmonella* spp. ed 1, p:33. Available at: <https://www.iso.org/standard/56714.html>.
- Joys TM (1985). The covalent structure of the phase-I flagellar filament protein of *Salmonella Typhimurium* and its comparison with other flagellins. The Journal of Biological Chemistry, 260:15758-15761. Available at: <https://www.jbc.org/content/260/29/15758>.
- Kim S, Frye JG, Hu J, Paula J, Cray F, Gautom R, and Boyle SD (2006). Multiplex PCR-based method for identification of common clinical serotypes of *Salmonella enterica* subsp. *enterica*. Journal of Clinical Microbiology, 44(10): 3608–3615. DOI:<https://doi.org/10.1128/JCM.00701-06>.
- Malorny B, Made D, Teufel P, Berghof-Jager C, Huber I, Anderson A and Helmuth R (2007). Multicenter validation study of two block cyclor- and one capillary-based real-time PCR methods for the detection of *Salmonella* in milk powder. International Journal of Food Microbiology, 117 (2): 211–218. DOI: <https://doi.org/10.1016/j.ijfoodmicro.2007.04.004>
- O'Regan E, McCabe E, Burgess C, McGuinness S, Barry T, Duffy G, Whyte P and Fanning S (2008). Development of a real-time multiplex PCR assay for the detection of multiple *Salmonella* serotypes in chicken samples. BMC Microbiology, 8: 156-167. DOI: <https://doi.org/10.1186/1471-2180-8-156>.
- Osman KM, Marouf SH, Erfan AM and Alattefy N (2014a). *Salmonella enterica* in imported and domestic day-old turkey poults in Egypt: Repertoire of virulence genes and their antimicrobial resistance profiles. OIE Revue Scientifique Technique, 33(3):1017-1026. DOI: <https://doi.org/10.20506/rst.33.3.2338>.
- Osman KM, Marouf SH, Zolnikov TR, and Alattefy N (2014b). Isolation and characterization of *Salmonella enterica* in day-old ducklings in Egypt. Pathogens and Global Health, 108(1):37-48. DOI: <https://doi.org/10.1179/2047773213Y.0000000118>.
- Pathmanathan SG, Cardona-Castro N, Sanchez-Jimenez MM, Correa-Ochoa MM, Puthuchery SD and Thong KL (2003). Simple and rapid detection of *Salmonella* strains by direct PCR amplification of the *hlyA* gene. Journal of Medical Microbiology, 52(9): 773–776. DOI: <https://doi.org/10.1099/jmm.0.05188-0>.
- Pereira F, Carneiro J, Matthiesen R, van Asch B, Pinto N, Gusmão L and Amorim A (2010). Identification of species by multiplex analysis of variable-length sequences. Nucleic Acids Research, 38 (22): 203. DOI:<https://doi.org/10.1093/nar/gkq865>. PM ID 20923781.
- Persad AK and LeJeune J (2018). A Review of Current Research and Knowledge Gaps in the Epidemiology of Shiga Toxin-Producing *Escherichia coli* and *Salmonella* spp. in Trinidad and

- Tobago. Veterinary Sciences, 5(2): 42. DOI:<https://doi.org/10.3390/vetsci5020042>.
- Prendergast DM, Hand D, NiGhallchóir E, McCabe, E, Fanning S, Griffin M, Egan J and Gutierrez M (2013). A multiplex real-time PCR assay for the identification and differentiation of *Salmonella enterica* serovar Typhimurium and monophasic serovar 4,[5],12:i:-. International Journal of Food Microbiology, 166(1): 48-53. DOI: <https://doi.org/10.1016/j.ijfoodmicro.2013.05.031>
- Sandjong BT, Sessitsch A, Liebana E, Kornschöber C, Allerberger F, Hächler H and Bodrossy L (2007). *Salmonella enterica* subsp. *enterica* serovar Typhimurium strain S05917 03 phase 1 flagellin (*fliC*) gene, partial cds. Journal of Microbiological Methods, 69(1): 23-36. Available at: [https://www.ncbi.nlm.nih.gov/gene?Db=gene&Cmd=Detail\\_sSearch&Term=1253480](https://www.ncbi.nlm.nih.gov/gene?Db=gene&Cmd=Detail_sSearch&Term=1253480).
- Shaw JA, Henard CA, Liu L, Dieckman LM, Andrés Vázquez-Torres AV and Bourret TJ (2018). *Salmonella enterica* serovar Typhimurium has three transketolase enzymes contributing to the pentose phosphate pathway. Journal of Biological Chemistry, 293(29): 11271-11282. DOI: <https://doi.org/10.1074/jbc.RA118.003661>.
- Shin HH, Hwang BH and Cha HJ (2016). Multiplex 16S rRNA-derived geno-biochip for detection of 16 bacterial pathogens from contaminated foods. Biotechnol Journal, 11(11): 1405-1414. DOI: <https://doi.org/10.1002/biot.201600043>.
- Silverman M, Zieg J and Hilmen M (1979). Phase variation in *Salmonella*: Genetic analysis of a recombinational switch. ResearchGate, 76(1): 391-5. DOI: <https://doi.org/10.1073/pnas.76.1.391>.
- Soumet C, Ermel G, Rose V, Drouin P, Salvat G and Colin P (1999). Identification by a multiplex PCR based assay of *Salmonella* Typhimurium and *Salmonella* Enteritidis strains from environmental swabs of poultry houses. Letters in Applied Microbiology, 29:1-6. DOI: <https://doi.org/10.1046/j.1365-2672.1999.00559.x>.
- Soyer Y, Moreno SA, Davis MA, Maurer J, McDonough PL, Schoonmaker-Bopp DJ, Dumas NB, Root T, Warnick LD, Gröhn YT and Wiedmann M (2009). *Salmonella enterica* serotype 4,5,12:i:-, an emerging *Salmonella* serotype that represents multiple distinct clones. Journal of Clinical Microbiology, 47: 3546-3556. DOI: <https://doi.org/10.1128/JCM.00546-09>
- Tamura K, Stecher G, Peterson D, Filipiński A and Kumar S (2013). MEGA6: molecular evolutionary genetics analysis version 6.0. Molecular Biology journal, 30: 2725-2729. DOI: <https://www.doi.org/10.1093/molbev/mst197>
- Tennant SM, Diallo S, Levy H, Livio S, Sow SO, Tapia M, Fields PI, Mikoleit M, Tamboura B, Kotloff KL, Nataro JP, Galen JE and Levine MM (2010). Identification by PCR of non-typhoidal *Salmonella enterica* serovars associated with invasive infections among febrile patients in Mali. PLoS Neglected Tropical Disease, 4(3): e621. DOI: <https://doi.org/10.1371/journal.pntd.0000621>.
- Thompson JD, Higgins DG and Gibson TJ (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids Research, 22(22): 4673-4680. DOI: <https://www.doi.org/10.1093/nar/22.22.4673>
- Woo PCY, Kenneth HLN, Susanna KPL, Kam-tong Yip, Ami MYF, Kitwah L, Dorothy MWT, Tak-lun Q, and Kwok-yung Y (2003). Usefulness of the MicroSeq 500 16S ribosomal DNA-based identification system for identification of clinically significant bacterial isolates with ambiguous biochemical profiles. Journal of Clinical Microbiology, 41: 1996-2001. DOI: <https://doi.org/10.1128/JCM.41.5.1996-2001.2003>.
- Yang X, Brisbin J, Yu H, Wang Q, Yin F, Zhang Y, Sabour P, Sharif S and Gong J (2014). Selected Lactic Acid-Producing Bacterial Isolates with the Capacity to Reduce *Salmonella* Translocation and Virulence Gene Expression in Chickens. PLOS ONE, 9(4): 93022. DOI: <https://doi.org/10.1371/journal.pone.0093022>