

# PARATYPHOID *SALMONELLA* SEROVARS IN CHICKENS: MOLECULAR DETECTION OF VIRULENCE AND ANTIMICROBIAL RESISTANCE GENES

Yousra M. NASSAR<sup>1</sup> , Wafaa A. ABD EL-GHANY<sup>1</sup>  , Adel K. IBRAHIM<sup>2</sup> , and Ahmed S. HAMOUDA<sup>1</sup> 

<sup>1</sup>Poultry Diseases Department, Faculty of Veterinary Medicine, Cairo University, Giza, 12211, Egypt

<sup>2</sup>Clinical Pathology Department, Faculty of Veterinary Medicine, Cairo University, Giza, 12211, Egypt

✉ Email: [wafaa.soliman@cu.edu.eg](mailto:wafaa.soliman@cu.edu.eg)

↳ Supporting Information

**ABSTRACT:** Paratyphoid salmonellosis is a serious disease that threatens the poultry industry worldwide, besides its public health hazard. The aims of this study were characterization of paratyphoid *Salmonella* spp. in chicken flocks of some Egyptian governorates, demonstration of the antimicrobial susceptibility of the isolated *Salmonella* spp., and detection of some virulence genes and antibiotic resistance genes using recent molecular techniques. A total of 238 organ samples were collected from 52 broiler, layer, and breeder chicken flocks, representing 9 Egyptian governorates. Conventional characterization of *Salmonella* isolates revealed a total isolation rate of 56.3% (134/238). Moreover, the isolation rates of *Salmonella* spp. were (49/79; 62%), (47/81; 58%), (10/18; 55.5%), (9/20; 45%), (2/6; 33.3%), (2/3; 66.7%), and (15/82; 53.6%) from liver, yolk sac, heart, spleen, caecum, ovary, and dead-in-shell embryos, respectively. A total of 32/238 (13.44%) isolates of *Salmonella* were found. Serological identification revealed presence of *S. enteritidis* (21.9%), *S. kentucky* (15.6%), *S. typhimurium* (12.5%), *S. molade* (12.5%), *S. takoradi* (9.4%), *S. wingrove* (6.3%), *S. infantis* (6.3%), *S. tsevie* (6.3%), *S. shangani* (3.1%), *S. bargny* (3.1%), and *S. papuana* (3.1%). All *Salmonella* strains (32/32; 100%) were resistant to streptomycin, while almost all of them (31/32; 96.9%) were susceptible to meropenem. The amplification of 16S rRNA gene of *Salmonella* isolates using uniplex polymerase chain reaction (PCR) generated a specific *Salmonella* product of approximately 550 base pair. The multiplex PCR revealed presence of *invA* (100%), *stn* (65.6%), and *sopB* (40.6%) virulence-associated genes as well as *aadA1* (100%), *blaTEM* (59.4%), *aadB* (18.75%), and *sul1* (28.1%) antibiotic resistance genes. In conclusion, virulent paratyphoid *Salmonella* spp. are circulating in the Egyptian flocks, causing economic losses. Additionally, they became resistant to the most commonly used field antibiotics. Therefore, regular molecular surveillance studies on the circulating *Salmonella* spp. and their resistance to the used antibiotics are of significant importance.

**Keywords:** Antibiotic resistance genes, Chicken, Paratyphoid *Salmonella*, PCR, Serology, Virulence

## INTRODUCTION

The poultry industry has grown dramatically around the world accounting for approximately 45% of global trade (Mottet and Tempio, 2017). This industry is considered as a significant source of animal protein for the Egyptians. Paratyphoid salmonellosis is one of the most important bacterial diseases affecting poultry and regarded a major source of food-borne infection in humans (Menghistu et al., 2011; Abd El-Ghany, 2020). Foodborne diseases are frequently associated with the consumption of animal-derived foods, primarily poultry products such as eggs and undercooked chicken (Castro-Vargas et al., 2020). These diseases have negative economic impacts on poultry industry as a result of the costs of investigation, surveillance, prevention, and treatment (Kuria, 2023). *Salmonella* spp. are rod-shape Gram-negative bacteria belonging to family *Enterobacteriaceae* and are known to infect many hosts (Xu et al., 2020).

Despite the European Union in 2006 prohibited using of some antibiotics as growth promoters feed additives, few countries are still using them (Gyles, 2008; Diab et al., 2019). Besides, many other antimicrobials are used haphazardly to control *Salmonella* infections either in poultry field or for human infection (Xu et al., 2020). As a result of indiscriminate use of these antibiotics, the emergence of multiple drug-resistant (MDR) *Salmonella* serotypes have significantly risen (Ammar et al., 2019). *Salmonella* spp. have developed resistance to a wide range of antimicrobials (Azizpour, 2018). The mechanisms by which the bacteria can acquire antibiotic resistance are variable and it can occur via mutations or horizontal transfer of resistance genes (Kapoor et al., 2017). However, MDR *Salmonella* spp. can be transmitted from poultry to humans through the food chain or via the direct contact (Firoozeh et al., 2012). The main threats affecting

**RESEARCH ARTICLE**  
 PII: S222877012200056-13  
 Received: June 23, 2023  
 Revised: September 24, 2023  
 Accepted: September 24, 2023

humans after exposure to these antibiotic-resistant bacteria are inability to treat patients successfully, along with the high risk of transmission of such resistant bacteria (Roca et al., 2015). Thus, the existence of antibiotic resistance *Salmonella* strains has become a serious global issue. In addition, presence of both antimicrobial resistance-associated genes as well as virulence associated genes in *Salmonella* strains are important indicators of pathogenicity degree (Capuano et al., 2013; Lamas et al., 2016). Among *Salmonella* spp., there are many virulence and antibiotic associated resistance genes that have been detected molecularly using polymerase chain reaction (PCR) assays. For instance, the virulence associated genes including *invA*, *sopB*, *stn*, and *pef* are used for genetic characterization of *Salmonella* strains (Badr et al., 2021; Elshebrawy et al., 2021; Ghetas et al., 2021; Hassan et al., 2021). Moreover, the antibiotic resistance genes such as *aadA1* (streptomycin), *aadB* (gentamicin), *sul1* (sulphonamides), and *bla*TEM ( $\beta$  lactams) have been detected in many isolates of *Salmonella* spp. (Ammar et al., 2019; Alam et al., 2020; Herrera-Sánchez et al., 2020).

Therefore, the aims of this study were characterization of paratyphoid *Salmonella* spp. in chicken flocks of some Egyptian governorates, demonstration of the antibiotic sensitivity of the isolated *Salmonella* spp., and detection of some virulence genes and antibiotic resistance genes using molecular techniques.

## MATERIALS AND METHODS

### Sampling

From November 2020 to April 2021, a total of 238 samples were collected from 52 broiler, layer, and breeder chickens' flocks in El-Mansoura, El-Sharqia, El-Qalyubia, El-Beheira, El-Minya, El-Fayoum, El-Gharbeya, Damietta, and Giza Egyptian governorates (Table 1). Samples were taken from diseased chicks showing whitish diarrhea, pasty vent, and omphalitis, dead-in-shell embryos, and breeders with low fertility and hatchability. The post-mortem lesions of both freshly dead and scarified diseased chickens were un-absorbed yolk sac, congested liver and spleen, typhlitis, and misshapen or discolored ova. Tissue samples were collected under aseptic conditions from liver (79), yolk sac (81), cecum (6), heart (18), spleen (20), ovary (3), gallbladders (3), and organs of dead-in-shell embryos (28), and then rapidly transferred in ice box to the laboratory for further processing.

### Ethical regulations

The study design and methodology were approved by the Institutional Animal Care and Use Committee (IACUC), Faculty of Veterinary Medicine, Cairo University, Egypt, with an approval number (01122022543).

### Isolation and identification

Isolation and identification of *Salmonella* spp. were done according to ISO 6579:2002 guidelines. The collected samples were inoculated in buffered peptone water and incubated at 37 °C for 18-20 hr. Then, 1 ml of pre-enriched broth was transferred into tubes containing 10 ml of Selenite F broth and incubated at 37±1 °C for 24 hr. A loopful from each broth culture was inoculated onto selective plating medium such as xylose lysine deoxycholate agar (XLD), brilliant green agar (BGA), and MacConkey agar media, incubated at 37 °C for 24 hr, and then checked for the of growth of typical *Salmonella* colonies. Suspected colonies were stained with Gram's and examined microscopically (Quinn et al., 2011). The biochemical identification of suspected *Salmonella* colonies was performed according to ISO 6579:2002 guidelines using triple sugar iron agar (TSI), indole, methyl red, vogues proskauer (VP), citrate utilization, urea hydrolysis, and H<sub>2</sub>S production tests. The serological identification of *Salmonella* was carried out according to Kauffman-White Scheme (Kauffman, 1974). Both somatic (O) and flagellar (H) antigens (DENKA SEIKEN Co., Japan) were demonstrated using polyvalent and monovalent O antisera according to a slide agglutination test and H antisera according to a tube agglutination test.

### The antimicrobial susceptibility test

The antimicrobial susceptibility test of the isolated *Salmonella* strains was done using disc diffusion method, according to the guidelines stipulated by National Committee for Clinical Laboratory Standards (NCCLS, 2020). The used antibiotic discs (Oxoid Limited, Basingstoke, Hampshire, UK) were ciprofloxacin (CP) (5µg), tetracycline (T) (30µg), ampicillin (AM) (10µg), cefotaxime (CF) (30µg), meropenem (M) (10µg), nalidixic acid (NA) (30µg), colistin (CO) (25µg), streptomycin (S) (10µg), levofloxacin (5µg), kanamycin (K) (30µg), clindamycin (CL) (10µg), amikacin (AK) (30µg), gentamicin (G) (10µg), and sulphamethoxazole (25µg). The tested strains were evaluated as susceptible, intermediate, and resistant by measuring the inhibitory zones. Moreover, the multiple antibiotic resistance (MAR) index for each *Salmonella* strain was determined (Singh et al., 2010) as follow;

MAR index= Number of antibiotics to which an isolate was resistant / total number of the tested antibiotics.

Each strain was classified as MDR if it exhibited resistance to three or more antimicrobial classes. Besides, the strain was considered to be resistant if it expressed MAR index > 0.2 (Shehata et al., 2019).

### Molecular detection of *Salmonella* spp. virulence-associated and antibiotic resistance genes

The detection of 16S rRNA of *Salmonella* isolates genes were carried out using uniplex PCR (Borges et al., 2017). The primers sequences are listed in Table 2, and the cycling conditions of primers during PCR are shown in Table 3. The

PCR reactions were adjusted in 25 µL reaction mixture containing 5 µL of DNA template, 12.5 µL of 2x PCR master mix, 1.25 µL each of forward and reverse primers (10 pmol/µL), and 5 µL nuclease free water. The PCR products were run through 1.5% agarose gel electrophoresis and a 100 bp DNA ladder was used as a size marker.

The multiplex PCR was used for testing the presence of both virulence associated genes and antibiotic resistance genes in the isolated *Salmonella* spp. The tested virulence associated genes were *invA*, *sopB*, *stn*, and *pef*, while the antibiotic resistance genes were *aadA1* (streptomycin), *aadB* (gentamicin), *sul1* (sulphonamides), and *blaTEM* (β lactams). The primers sequences of virulence associated genes and the antibiotic resistance genes are presented in Tables 4 and 5, respectively.

## RESULTS

Suspected *Salmonella* isolates appeared as small, non-lactose fermenter, colorless, and transparent colonies on MacConkey agar plates. However, isolates appeared as pink colonies with black centers on XLD and pinkish white or red colonies surrounded by a red halo on BGA agar. Gram staining revealed presence of Gram-negative, medium size, and rod-shaped bacilli under the microscope. The isolates fermented glucose, but did not ferment lactose and sucrose on TSI slants, and appeared as red slant with yellow butt, with or without H<sub>2</sub>S production. Moreover, they were negative for indole, VP, and urea hydrolysis, whereas positive for methyl red and citrate utilization.

The conventional bacteriological characterization revealed that the isolation rates of *Salmonellae* were (49/79; 62%), (47/81; 58%), (10/18; 55.5%), (9/20; 45%), (2/6; 33.3%), (2/3; 66.7%), and (15/82; 53.6%) from liver, yolk sac, heart, spleen, caecum, ovaries, and dead-in-shell embryos, respectively. However, no isolation (0/3; 0%) was obtained from gallbladder (Table 6). The total isolation rate was 134/238 (56.3%). The number of the total strains of *Salmonella* was 32/238 (13.44%). Moreover, positive *Salmonella* isolates were detected in 49 out of 76 in broilers, 4 out of 10 in layers, and 2 out of 3 in breeder chicken flocks.

By using polyvalent and monovalent O and H antiserum, the slide and tube agglutination tests confirmed that the tested *Salmonella* isolates (n= 32) belonged to 11 paratyphoid *Salmonella* serovars. Table 7 shows the types and percentages of the isolated *Salmonella* serovars. A total of 11 serovars of paratyphoid *Salmonella* were identified as *S. enteritidis* (7, 21.9%), *S. kentucky* (5, 15.6%), *S. typhimurium* (4, 12.5%), *S. molade* (4, 12.5%), *S. takoradi* (3, 9.4%), *S. wingrove* (2, 6.3%), *S. infantis* (2, 6.3%), *S. tsevie* (2, 6.3%), *S. shangani* (1, 3.1%), *S. bargny* (1, 3.1%), and *S. papuana* (1, 3.1%).

Paratyphoid *Salmonella* serovars isolated from the liver (n= 49) were *S. enteritidis* (12), *S. typhimurium* (6), *S. kentucky* (8), *S. takoradi* (7), *S. molade* (4), *S. wingrove* (3), *S. bargny* (3), *S. shangani* (1), *S. infantis* (2), and *S. tsevie* (3), while those isolated from the yolk sac (n= 47) were *S. enteritidis* (9), *S. typhimurium* (4), *S. kentucky* (9), *S. takoradi* (6), *S. molade* (7), *S. wingrove* (2), *S. bargny*(3), *S. shangani* (1), *S. infantis* (2), and *S. tsevie* (4). Moreover, strains of the heart (n= 10) were *S. enteritidis* (2), *S. typhimurium* (3), *S. wingrove* (1), *S. shangani* (1), *S. molade* (2), and *S. infantis* (1), but those of spleen (n= 9) were *S. enteritidis* (1), *S. typhimurium* (5), *S. kentucky* (1), *S. papuana* (1), and *S. wingrove* (1). Caecal serovars (n= 2) were *S. takoradi* (2), while ovarian serovars (n= 2) were *S. enteritidis* (1) and *S. molade* (1). Regarding serovars of the dead-in-shell embryos (n= 15), they were *S. kentucky* (3), *S. takoradi* (6), *S. molade* (3), and *S. bargny* (3).

Regarding the distribution of *Salmonella* serovars in the surveyed Egyptian governorates, the highest isolation rates were from El-Dakhliya, El-Sharqia, Damietta and El-Qalubia, followed by El-Fayoum, El-Minya, Giza, and El-Beheira. Nevertheless, there was no isolation from El-Gharbeya. The data in Table 8 reveals detection of *S. enteritidis* from El-Qalubia, El-Dakhliya, El-Beheira, El-Minya, El-Fayoum, and Damietta governorates, while *S. typhimurium* was found in El-Dakhliya, El-Fayoum, El-Sharqia, and Giza. Moreover, *S. kentucky* was isolated from El-Dakhliya, El-Minya, and Damietta and *S. molade* from El-Beheira, El-Fayoum, and Damietta. *S. takoradi* was isolated from El-Dakhliya, El-Sharqia, and Damietta, though *S. wingrove* was demonstrated in El-Dakhliya and El-Qalubia. Both *S. shangani* and *S. bargny* were detected in El-Dakhliya, then *S. tsevie* was isolated from El-Dakhliya and El-Sharqia. *S. infantis* was demonstrated in El-Fayoum and El-Sharqia, whereas *S. Papuana* was found in El-Qalubia

As shown in Table 9, all strains of *Salmonella* were completely resistant to streptomycin 100% (32/32), followed by clindamycin 93.8% (30/32), nalidixic acid 75% (24/32), amikacin 65.6% (21/32), tetracycline 50% (16/32), cefotaxime 43.7% (14/32), kanamycin 34.4% (11/32), sulphamethoxazole 28.1% (9/32), ampicillin 18.8% (6/32), ciprofloxacin 12.5% (4/32), and colistin 12.5% (4/32). However, almost all strains of *Salmonella* were susceptible to meropenem 96.9% (31/32), followed by levofloxacin 90.6% (29/32), gentamicin 87.5% (28/32), colistin 84.4% (27/32), ciprofloxacin 81.3% (26/32), ampicillin 68.8% (22/32), sulphamethoxazole 71.9% (23/32), kanamycin 59.3% (19/32), and cefotaxime 56.3% (18/32).

Additionally, a total of 24 out of 32 *Salmonella* strains were resistant to 3 or more of the tested antimicrobials with an average MAR index 0.350. Surprisingly, out of the tested (n= 14) antimicrobials, two strains of *S. enteritidis* were resistant to 14, one strain of *S. kentucky* was resistant to 13 antimicrobials, and one strain of *S. typhimurium* was resistant to 10 antimicrobials.

The results of uniplex PCR confirmed that the amplification of 16S rRNA gene of all *Salmonella* isolates generated a product of approximate molecular size of 550 base pair (bp) (Figure 1). Concerning the virulence genes, the amplified

products for *invA*, *sopB*, *stn*, and *pef* genes were 284, 517, 617, and 700 bp, respectively, while those of the antibiotic resistance genes *aadA1*, *aadB*, *sul1*, and *blaTEM* were 629, 300, 591, and 608 bp, respectively.

All *Salmonella* strains (n= 32) were screened by multiplex PCR for identification of the virulence associated genes (*invA*, *stn*, *sopB*, and *pef*) and the antibiotic resistance genes (*aadA1*, *aadB*, *sul1*, and *blaTEM*). The results indicated that *invA*, *sopB*, and *stn* genes were harbored by 100%, 40.6%, and 65.6% of *Salmonella* strains, respectively, while none of them had *pef* gene (Table 10 and Figure 2). Further, all strains (100%) harbored *aadA1* associated with streptomycin resistance, but 59.4%, 18.75%, and 6.3% of strains possessed *blaTEM* (ampicillin), *aadB* (gentamicin), and *sul1* (sulphamethoxazole) resistance genes, respectively (Table 11 and Figures 3-5).

**Table 1 - The source and number of the samples examined for detection of *Salmonella***

Governorates	Chickens			Dead-In-shell embryonic organs	No. of samples	No. of flocks
	Broiler	Layer	Breeder			
A	39	6	3	28	145	28
B	18	2	0	0	49	14
C	19	2	0	0	44	10
Total	76	10	3	28	238	52

A= El-Mansoura, El-Gharbeya, El-Beheira & Damietta; B= El-Qalyubia, El-Sharqia & Giza; C= El-Minya & El-Fayoum

**Table 2 - Primer sequences for detection of 16S rRNA gene of *Salmonellae***

Target gene	Primer Sequence	Gene	Reference
16S rRNA	GCA ACG CGA AGA ACC TTA CC (Forward) GGT TAC CTT GTT ACG ACT T (Reverse)	16S rRNA	Gopinath et al. (1998)

**Table 3 - Cycling conditions of the used primers during PCR**

Gene	Primary Denaturation	Secondary Denaturation	Annealing	Extension	No. of Cycles	Final Extension
16S rRNA gene and other genes	90 °C for 5 min.	90 °C for 1 min.	52 °C for 1 min.	72 °C for 1 min.	39	72 °C for 7 min.

**Table 4 - Primers sequences of virulence associated genes of *Salmonella* serovars**

Target gene	Primer sequence	Virulence factor	Reference
<i>invA</i>	GTGAA ATT ATC GCC ACG TTC GGG CAA (Forward) TCAT CGC ACC GTC AAA GGA ACC (Reverse)	<i>Salmonella</i> species/SPI-1	Oliveira et al. (2003)
<i>stn</i>	TTG TGT CGC TAT CAC TGG CAA CC (Forward) ATT CGT AAC CCG CTC TCG TCC (Reverse)	Enterotoxin/Chromosome	Murugkar et al. (2003)
<i>sopB</i>	TCA GAA GRC GTC TAA CCA CTC (Forward) TAC CGT CCT CAT GCA CAC TC (Reverse)	Translocated effector protein/SPI-5	Huehn et al. (2010)
<i>pef</i>	TGT TTC CGG GCT TGT GCT (Forward) CAG GGC ATT TGC TGA TTC C (Reverse)	Plasmid encoded fimbriae/Plasmid	Murugkar et al. (2003)

**Table 5 - Primers sequences of antiotic resistance genes of *Salmonella* serovars**

The antibiotic	Target gene	Primer sequence	Reference
Streptomycin	aadA1	CTC CGC AGT GGA TGG CGG (Forward)	Chuanchuen and Padungtod (2009)
		GAT CTG CGC GCG AGG CCA (Reverse)	
Gentamicin	aadB	CTAGCTGCGGCAGATGAGC (Forward)	
		CTCAGCCGCTCTGGGCA (Reverse)	
Sulfamethoxazole	sul1	CGGACGCGAGGCCTGTATC (Forward)	
		GGGTGCGGACGTAGTCAGC (Reverse)	
Ampicillin	blaTEM	ATCAGTTGGGTGCACGAGTG (Forward)	
		ACGCTCACCGCTCCAGA (Reverse)	

**Table 6 - The number and percentage of the positive *Salmonella* samples**

Group	Liver		Yolk sac		Cecum		Heart		Spleen		Gallbladder		Ovary		Dead-In-shell embryos		Total	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
A	26/47	55.3	26/51	50.9	2/5	40	1/1	100	3/10	30	0	0	2/3	66.6	15/28	53.6	75/145	51.7
B	12/16	75	10/14	71.4	0/1	0	5/9	55.5	5/6	83.3	0/3	0	0	0	0	0	32/49	56.3
C	11/16	68.8	11/16	68.8	0	0	4/8	50	1/4	25	0	0	0	0	0	0	27/44	61.4
Total	49/79	62	47/81	58	2/6	33.3	10/18	55.5	9/20	45	0/3	0	2/3	66.7	15/28	53.6	134/238	56.3

A= El-Mansoura, El-Gharbeya, El-Beheira & Damietta; B= El-Qalyubia, El-Sharqia & Giza; C= El-Minya & El-Fayoum

**Table 7 - The types and percentages of the isolated *Salmonella* serovars**

Identified strains	Number	%	Antigenic structure		
			Group	O	H
<i>S. enteritidis</i>	7	21.9	D1	1,9,12	g,m
<i>S. kentucky</i>	5	15.6	C3	8,20	i : Z6
<i>S. typhimurium</i>	4	12.5	B	1,4,5,12	i : 1,2
<i>S. molade</i>	4	12.5	C2	8,20	Z10 : Z6
<i>S. takoradi</i>	3	9.4	C2	8,20	i : 1,5
<i>S. wingrove</i>	2	6.3	C2	6,8	c : 1,2
<i>S. infantis</i>	2	6.3	C1	6,7	r : 1,5
<i>S. tsevie</i>	2	6.3	B	4,5	i : e,n,z15
<i>S. shangani</i>	1	3.1	E1	3,10	d : 1,5
<i>S. bargny</i>	1	3.1	C3	8,20	i : 1,5
<i>S. papuana</i>	1	3.1	C1	6,7	r : e,n,Z15

**Table 8 - Types of *Salmonella* serovars isolated from the Egyptian governorates**

<i>Salmonella</i> serovar	Frequency of each isolate		Governorate
	No.	%	
<i>S. enteritidis</i>	7	21.9	El-Qalubia, El-Dakhliya, El-Beheira, El-Minya, El-Fayoum, & Damietta
<i>S. kentucky</i>	5	15.6	El-Dakhliya, El-Minya, & Damietta
<i>S. typhimurium</i>	4	12.5	El-Dakhliya, E-Fayoum, El-Sharqia, & Giza
<i>S. molade</i>	4	12.5	El-Beheira, El-Fayoum, & Damietta
<i>S. takoradi</i>	3	9.4	El-Dakhliya, El-Sharqia, & Damietta
<i>S. wingrove</i>	2	6.3	El-Dakhliya and El-Qalubia.
<i>S. infantis</i>	2	6.3	El-Fayoum & El-Sharqia.
<i>S. tsevie</i>	2	6.3	El-Dakhliya & El-Sharqia
<i>S. shangani</i>	1	3.1	El-Dakhliya
<i>S. bargny</i>	1	3.1	El-Dakhliya
<i>S. papuana</i>	1	3.1	El-Qalubia

**Table 9 - The antimicrobial susceptibility test of the isolated *Salmonella* strains**

<i>Salmonella</i> serovars	No.	Antibiotic resistance													
		S	CL	NA	AK	T	CF	K	SXT	AM	CP	CO	G	L	M
<i>S. enteritidis</i>	7	100	100	85.7	85.7	71.4	57.1	42.9	42.9	28.5	28.5	28.5	14.3	4.3	4.3
<i>S. kentucky</i>	5	100	80	80	60	60	60	60	40	40	20	20	20	20	0
<i>S.typhimurum</i>	4	100	100	75	50	50	25	50	25	25	0	25	0	0	0
<i>S. molade</i>	4	100	100	75	75	50	50	25	25	25	25	0	0	0	0
<i>S. takoradi</i>	3	100	100	66.6	66.6	33.3	33.3	33.3	33.3	0	0	0	0	0	0
<i>S. wingrove</i>	2	100	100	100	50	50	50	50	50	0	0	0	0	0	0
<i>S. infantis</i>	2	100	100	50	50	50	50	0	0	0	0	0	0	0	0
<i>S. tsevie</i>	2	100	50	50	50	50	50	0	0	0	0	0	0	0	0
<i>S. shangani</i>	1	100	100	100	100	0	0	0	0	0	0	0	0	0	0
<i>S. bargny</i>	1	100	100	100	100	0	0	0	0	0	0	0	0	0	0
<i>S. papuana</i>	1	100	100	0	0	0	0	0	0	0	0	0	0	0	0
Total	32	32	30	24	21	16	14	11	9	6	5	4	2	2	0
Resistant %		100	93.8	75	65.6	50	43.7	34.4	28.1	18.8	12.5	12.5	6.3	6.3	3.1
Intermediate %		0	3.1	9.4	9.4	6.3	0	6.3	3.1	12.5	6.3	3.1	6.3	3.1	0
Susceptible %		0	3.1	15.6	25	43.7	56.3	59.3	68.8	68.8	78.1	81.3	87.5	90.6	96.9

S: Streptomycin; CL: Clindamycin; NA: Nalidixic acid; AK: Amikacin; T: Tetracycline; CF: Cefotaxim; K: Kanamycin; SXT: Sulphamethoxazol; AM: Ampicillin; CP: Ciprofloxacin; CO: Colistin; G: Gentamicin; L: Levofloxacin; M: Meropenem


**Table 10 - Percentages of virulence-associated genes of *Salmonella* serovars**

Virulence genes <i>Salmonella</i> serovar	No.	16S rRNA		<i>invA</i>		<i>stn</i>		<i>sopB</i>		<i>pef</i>	
		No.	%	No.	%	No.	%	No.	%	No.	%
<i>S. enteritidis</i>	7	7	100	7	100	6	85.7	5	71.5	0	0
<i>S. kentucky</i>	5	5	100	5	100	4	80	2	40	0	0
<i>S. typhimurum</i>	4	4	100	4	100	4	100	1	25	0	0
<i>S. molade</i>	4	4	100	4	100	1	25	0	0	0	0
<i>S. takoradi</i>	3	3	100	3	100	1	33.3	1	33.3	0	0
<i>S. wingrove</i>	2	2	100	2	100	2	100	2	100	0	0
<i>S. infantis</i>	2	2	100	2	100	1	50	1	50	0	0
<i>S. tsevie</i>	2	2	100	2	100	1	50	1	50	0	0
<i>S. shangani</i>	1	1	100	1	100	1	100	0	0	0	0
<i>S. bargny</i>	1	1	100	1	100	0	0	0	0	0	0
<i>S. papuana</i>	1	1	100	1	100	0	0	0	0	0	0
Total	32	32	100	32	100	21	65.6	13	40.6	0	0


**Table 11 - Percentages of antibiotic associated genes of *Salmonella* serovars**

<i>Salmonella</i> serovar	No.	The tested antibiotic resistance genes							
		<i>aadA1</i>		<i>sul1</i>		<i>blaTEM</i>		<i>aadB</i>	
		No.	%	No.	%	No.	%	No.	%
<i>S. enteritidis</i>	7	7	100	3	42.8	3	42.8	0	0
<i>S. kentucky</i>	5	5	100	2	40	3	60	2	40
<i>S. typhimurum</i>	4	4	100	1	25	2	50	0	0
<i>S. molade</i>	4	4	100	1	25	2	50	1	25
<i>S. takoradi</i>	3	3	100	1	33.3	1	33.3	1	33.3
<i>S. wingrove</i>	2	2	100	1	33.3	2	100	0	0
<i>S. infantis</i>	2	2	100	0	0	2	100	1	50
<i>S. tsevie</i>	2	2	100	0	0	1	50	1	50
<i>S. shangani</i>	1	1	100	0	0	1	100	0	0
<i>S. bargny</i>	1	1	100	0	0	1	100	0	0
<i>S. papuana</i>	1	1	100	0	0	1	100	0	0
Total	32	32	100	9	28.1	19	59.4	6	18.75


**Figure 1 - Agarose gel electrophoresis for amplified samples, where Lanes 1, 2, 3, 4, 7, 8, and 9 represent the positive *Salmonella* strains, Lane 5: Positive control showing 550 bp for the amplified 16S rRNA gene, Lane M: 100-bp DNA Marker and Lane 6: Negative control**



**Figure 2** - Multiplex PCR products of *Salmonella* virulence genes. Lane 1: Positive control, Lanes 2, 5, 6, 7, and 8 showing amplicons of 284 bp, 517 bp, and 671 bp of *invA*, *sopB* and *stn* genes respectively, Lanes 3 and 4 showing amplicons of 284 bp of *invA* gene and Lane M: 100-bp DNA marker



**Figure 3** - Agarose gel electrophoresis for PCR products of antibiotic resistance gene (streptomycin) of *Salmonella* showing amplicon of 629 bp of *aadA1* gene. Lane 1: Negative Control, Lane M: 100-bp DNA marker, and Lane 2: Positive control



**Figure 4** - Agarose gel electrophoresis for PCR products of antibiotic resistance gene (sulfamethoxazole) of *Salmonella* showing amplicon of 591 bp of *sul1* gene. Lane M: 100-bp DNA marker, Lane 3: Positive control and Lane 4: Negative Control






Figure 5 - Multiplex PCR products of antibiotic resistance genes (ampicillin and gentamicin) of *Salmonella* showing amplicon of 608 bp of *bla*<sub>TEM</sub> gene and 300bp of *aadB* gene, respectively. Lane 1: Negative Control, Lane M: 100-bp DNA marker and Lane 2: Positive control.

## DISCUSSION

Paratyphoid salmonellosis is a prevalent foodborne disease which has a global economic and public health concern (Aziz et al., 2018). Therefore, this study characterized paratyphoid *Salmonella* spp. in chicken flocks of some Egyptian governorates. Our results indicated that *S. enteritidis* was the most prevalent serovar (21.8%), followed by *S. kentucky* (15.6%) and *S. typhimurium* (12.5 %). These results are in agreement with the findings of Elsayed et al. (2019) who detected *S. enteritidis* (38.8%), *S. kentucky* (23.3%), and *S. typhimurium* (23.3%). Further, Elshebrawy et al. (2021) detected similar observation. Many reports indicated that *S. enteritidis* is one of the predominant circulating serovars in poultry flocks, as it has been isolated in rates of 58.33% (Rabie et al., 2012), 55.6% (Moussa et al., 2010), and 37.25% (Abd El-Ghany et al., 2012). Additionally, *S. typhimurium* is another most frequently reported and prevalent serovar worldwide (WHO, 2006). The study of Ammar et al. (2016) indicated that the isolation rates of *S. typhimurium* and *S. enteritidis* were 52.94% and 11.76%, respectively (Ammar et al., 2016) and 86.6% and 9%, respectively (El-Sharkawy et al., 2017). Regarding the low isolation rates of *S. tsevie* (6.3%) and *S. papuana* (3.1%) in this study, nearly similar results were obtained by Al-baqir et al. (2019) and Abd El-Tawab et al. (2015) who reported that the isolation rate of *S. tsevie* was (2%) and *S. papuana* was 2.3%, respectively. Other serovars of *Salmonella* such as *S. wingrove* (6.3%) and *S. shangani* (3.1%) were detected here, which may be due to improper biosecurity measures within farms and the possibility of disease transmission via various reservoirs and farm workers (Elshebrawy et al., 2021).

The highest isolation rate of *Salmonella* serovars was from the liver (62%), rather than the other tissues examined. The previous reports of Menghistu et al. (2011), Abdel-Aziz (2016), Al-baqir et al. (2019), and Saleem et al. (2022) mentioned similar finding. It is important to note that the high detection of *Salmonella* in the liver live may indicate the potential of the pathogen to cause a systemic infection.

It has been noticed that the highest isolation of *Salmonella* serovars was from El-Dakhliya, El-Sharqia Damietta, and El-Qalubia followed by El-Fayoum, El-Minya, Giza, and El-Beheira governorate. These results are in accordance with the fact that the highest percentages of chicken farms allocated in El-Dakhliya, El-Sharqia, Damietta and El-Qalubia rather than other Egyptian governorates.

The development of MDR *Salmonella* strains has emerged as a major public health concern around the world (Marshall and Levy, 2011). This resistance poses a direct threat to human health when treatment is hindered without a complete course and when there are interactions with both animals and humans pathogens (Frye and Jackson, 2013).

Concerning the antimicrobial susceptibility test, the results of the current study indicated that all of the isolated *Salmonellae* strains were resistant to streptomycin (100%), which may be due to the misuse of this antibiotic in the poultry field. However, other studies reported that 100% (Habrun et al., 2012) or 89.7% (El-Sharkawy et al., 2017) of isolates were sensitive to streptomycin.

On the other side site, 96.9% of the isolated *Salmonella* strains were susceptible to meropenem. The same result was obtained by Abou Elez et al. (2021). Furthermore, Elshebrawy et al. (2021) detected a low resistance rate (7.6%) of *Salmonella enterica* serovars against meropenem. It has been reported that carbapenems could be used in the treatment

of salmonellosis in MDR cases (Calayag et al., 2017). The low resistance rate against meropenem could be attributed to the underuse of carbapenems in veterinary medicine particularly for chicken infections.

Here, the resistance of the isolates to ciprofloxacin was 15.6%, which agrees with that of Donado-Godoy et al. (2012) (15%). In comparison, the resistance to ciprofloxacin was 50% (Abdel Rahman et al., 2014), 35% (Al-baqir et al., 2019), 33% (Badr et al., 2021), or 27.8% (Elshebrawy et al., 2021). As well, the resistance to ampicillin was 18.8%, which was much less than Diab et al. (2019) (68.2%), Ghetas et al. (2021) (78.5%), Alam et al. (2020) (82.85%), Nabil and Yonis, (2019) (94.1%), and Raji et al. (2021) (100%). Further, 50% of the isolated *Salmonellae* showed resistance to tetracycline, which nearly agreed, but lower than those stated by Raji et al. (2021) (75%) and Ibrahim et al. (2021) (62%). The maximum resistance to tetracycline (97.14%) was reported by Alam et al. (2020).

Despite conventional cultural methods offer the advantage of being able to detect live cells and evaluate large numbers of samples, they are time-consuming and laboring (Maciorowski et al., 2006; Margot et al., 2013). On the other hand, molecular techniques such as PCR are rapid, specific, and sensitive, and could replace traditional detection methods (Siddique et al., 2009; Ibrahim et al., 2014). Moreover, multiplex PCR platform provides a greater detection efficiency, allows simultaneous detection of several diseases at the same time, and saves both money and time (Lee et al., 2014; Li et al., 2020).

In the current study, the amplification of 16S rRNA gene of all *Salmonella* isolates gave a unique specific product at 550 bp. Also, the amplified products of the virulence genes (*invA*, *sopB*, *stn*, and *pef*) were 284, 517, 617, and 700 bp, respectively. The *invA* protein is an inner membrane component of the *Salmonella* pathogenicity island 1 type 3 secretion system (Shah et al., 2011). The *invA* gene allows bacteria to invade the host cells (Cha et al., 2013). This gene is only found in *Salmonella* spp. and is therefore a valuable diagnostic tool for genetic identification (O'Regan et al., 2008). Several studies have noted a high frequency of *invA* virulence gene in *Salmonella* serovars (Karmi, 2013). All examined *Salmonella* serovars (100%) contained *invA* gene. Many other studies agreed with ours (Shabnam and Kwai, 2010; Campioni et al., 2012; Samanta et al., 2014; Radwan et al., 2016; Amini et al., 2018; Hassan et al., 2018; ElSheikh et al., 2019; Ramatla et al., 2020; Elshebrawy et al., 2021).

The *Salmonella* enterotoxin (*stn*) gene encodes the Stn protein which causes gastroenteritis and regulates the integrity of bacterial cell membranes (Huehn et al., 2010). This gene was found in 65.6% of the strains in this study, which is similar to Elshebrawy et al. (2021) (65.8%). Other researchers have also noted high rates of *stn* gene (Zou et al., 2012; Osman et al., 2014; Ammar et al., 2019; Sabry et al., 2020; Hassan et al., 2021).

Fimbriae are essential for *Salmonella* pathogenicity, as they facilitate the attachment of the pathogen to epithelial cells. The *pef* gene encodes *pef* fimbria (Murugkar et al., 2003; Ammar et al., 2016). The examined isolates in the present study had no *pef* gene, which is similar to the results reported by Elkenany et al. (2019), Elhariri et al. (2020), and Hassan et al. (2021). Also, a low frequency of the *pef* gene was detected by Ahmed et al. (2016) (6.7%) and Ghetas et al. (2021) (21%). This could be explained by the fact that bacteria use other fimbriae to attach to the host cells. However, in other studies, various findings were noted; for example, Awad et al. (2020) identified *pef* gene in 54.8% of the isolates and highlighted the role of fimbriae in the infection.

By triggering secretory pathways, promoting inflammation or changing ion balances within cells, the *sopB* gene contributes to the development of diarrhea (Ahmed et al. 2016). The prevalence of the *sopB* gene is 40.6% which is lower than Awad et al. (2020) (87.1%), Mohamed et al. (2021) (91.3%), and Badr et al. (2021) (100%).

The amplified products of the antibiotic resistance genes (*aadA1*, *aadB*, *sul1*, and *blaTEM*) were 629, 300, 591, and 608 bp, respectively. In this study, we observed that the phenotypical resistance to ampicillin is 18.8%, but 59.4% of the isolates tested positive for the *blaTEM* gene by PCR, a gene encoding for resistance to  $\beta$ -lactams. This assumes that some antimicrobial resistance genes in some bacteria are inactive or "silent" *in-vitro*. However, these silent genes can spread to other bacteria or can become active *in-vivo* under antimicrobial pressure (Ma et al., 2007). In contrast, El-Sharkawy et al. (2017) found that all isolated strains of *S. enteritidis* were ampicillin-susceptible and *blaTEM* negative, suggesting that these strains had a different ampicillin resistance mechanism. In a previous work, according to Aslam et al. (2012), the *blaTEM* gene was present in 17% of *Salmonella* spp. isolated from retail meats in Canada and was the most frequently found resistance gene. According to Lu et al. (2011), 81.2 % of the 108 *S. indiana* isolates tested positive for the *blaTEM* gene. Ammar et al. (2019) noticed the *blaTEM* gene in 100% of the isolates. The genes *blaPSE-1* and *blaTEM*, which encode  $\beta$ -lactamases conferring resistance to ampicillin, were found in 69.4% of the strains, according to a report by Herrera-Sánchez et al. (2020). In the present study, we found out that all isolates of *Salmonella* were completely resistant to streptomycin and the gene for streptomycin resistance, *aadA1* was found in all the isolated strains. This rate is higher than those reported by Chuanchuen and Padungtod, (2009), who detected the resistance gene *aadA1* in (17%), Doosti et al. (2017) (45.6%), Alam et al. (2020) (77.1%), and Herrera-Sánchez et al. (2020) (87.8%).

6.3% of the isolated *Salmonella* strains were resistant to gentamycin, however the *aadB* gene that confers resistance to gentamycin was found in 18.75% of the isolated strains, as the gene was silenced. In contrast, the *aadB* gene was not found in any of the strains in the study conducted by Herrera-Sánchez et al. (2020).

A percentage of 28.1 of the isolated strains were phenotypically resistant to sulphamethoxazole and 28.1% of isolates being positive for *sul1* which confer sulphamethoxazole resistance. This result is lower than those reported by Adesiji et al. (2014) (100%), Mohamed and Suelam (2010) (97.3%), Aziz et al. (2018) (83.3%), and El-Sharkawy et al. (2017) (57%).

## CONCLUSION

Paratyphoid *Salmonella* spp. found in poultry flocks at high rates pose a zoonotic danger. Additionally, MDR *Salmonella* serovars with a diverse range of virulence genes have been detected in the existed *Salmonella* spp. These results indicate the importance of the constant surveillance of antibiotic resistant *Salmonella* strains, the use of alternatives instead of antimicrobials in poultry, and adoption of strong public health and food safety protocols to reduce the human health risk associated with salmonellosis.

## DECLARATIONS

### Corresponding author

E-mail: wafaa.soliman@cu.edu.eg

### Author's contribution

This work was carried out with the contribution of all authors. Nassar YM conducted the practical part of the research and helped in data collection, analysis, and writing. Ibrahim AA conducted the molecular work and validated it. Hamouda AS, Abd El-Ghany WA, and Ibrahim AA designed the protocol, supervise the work, and approved it. Abd El-Ghany WA helped in data collection and write, formatted, and submit the draft of the manuscript. All authors read and approved the final version of the manuscript.

### Conflict of interest

The authors declare that they have no conflicts of interest.

## REFERENCES

- Abd El-Ghany W A (2020). Salmonellosis: A food borne zoonotic and public health disease in Egypt. *Journal of infection in the developing Countries*, 14(7): 674-678. DOI: <https://doi.org/10.3855/jidc.12739>
- Abd El-Ghany WA, El-Shafii SS, and Hatem ME (2012). A survey on *Salmonella* species isolated from chicken flocks in Egypt. *Asian Journal of Animal and Veterinary Advances*, 7(6): 489-501. DOI: <https://doi.org/10.3923/ajava.2012.489.501>
- Abd El-Tawab A, Ashraf A, El-Hofy FI, Ammar AM, Nasef SA, and Nabil NM (2015). Studies on different *Salmonella* serotypes isolated from poultry in different governorates in Egypt. *Benha Veterinary Medical Journal*, 28(2): 169-175. DOI: <https://doi.org/10.21608/bvmj.2015.32498>
- Abdel Rahman M, Hamed E, Roshdy H, Sorour H, and AlAtfeehy N (2014). Bacteriological studies on native and imported apparently healthy one day old chicks. *Suez Canal Veterinary Medical Journal*, 19(2): 263-276. DOI: <https://dx.doi.org/10.21608/scvmj.2014.65854>
- Abdel-Aziz NM (2016). Detection of *Salmonella* species in chicken carcasses using genus specific primer belong to *invA* gene in Sohag city, Egypt. *Veterinary World*, 9(10): 1125-1128. DOI: <https://doi.org/10.14202/vetworld.2016.1125-1128>
- Abou Elez RM, Elsohaby I, El-Gazzar N, Tolba HM, Abdelfatah EN, Abdellatif SS, et al. (2021). Antimicrobial resistance of *Salmonella enteritidis* and *Salmonella typhimurium* isolated from laying hens, table eggs, and humans with respect to antimicrobial activity of biosynthesized silver nanoparticles. *Animals*, 11(12): 3554. DOI: <https://doi.org/10.3390%2Fani11123554>
- Adesiji YO, Deekshit VK, and Karunasagar I (2014). Antimicrobial-resistant genes associated with *Salmonella* spp. isolated from human, poultry, and seafood sources. *Food Science & Nutrition*, 2(4): 436-442. DOI: <https://doi.org/10.1002/fsn3.119>
- Ahmed HA, El-Hofy FI, Shafik SM, Abdelrahman MA, and Elsaid GA (2016). Characterization of virulence-associated genes, antimicrobial resistance genes, and class 1 integrons in *Salmonella enterica* serovar Typhimurium isolates from chicken meat and humans in Egypt. *Foodborne Pathogens and Disease*, 13(6): 281-288. DOI: <https://doi.org/10.1089/fpd.2015.2097>
- Alam SB, Mahmud M, Akter R, Hasan M, Sobur A, Nazir KNH, et al. (2020). Molecular detection of multidrug resistant *Salmonella* species isolated from broiler farm in Bangladesh. *Pathogens*, 9(3): 201. DOI: <https://doi.org/10.3390/pathogens9030201>
- Al-baqir A, Hussein A, Ghanem I, and Megahed M (2019). Characterization of paratyphoid *Salmonellae* isolated from broiler chickens at Sharkia governorate, Egypt. *Zagazig Veterinary Journal*, 47(2): 183-192. DOI: <https://doi.org/10.21608/zvzj.2019.10544.1028>
- Amini K, Salehi TZ, Nazari Z, Nikbakht G, Ranjbar R, Amini J, et al. (2018). Molecular detection of *invA* and *spv* virulence genes in *Salmonella typhimurium* isolated from human and animals in Iran. *Advanced Journal of Microbiology Research*, 12 (3): 001-009. Available at: <https://www.international-scholars-journals.com/articles/molecular-detection-of-inva-and-spv-virulence-genes-in-salmonella-enteritidis-isolated-from-human-and-animals-in-iran.pdf>
- Ammar AM, Abdeen EE, Abo-Shama UH, Fekry E, and Kotb Elmahallawy E (2019). Molecular characterization of virulence and antibiotic resistance genes among *Salmonella* serovars isolated from broilers in Egypt. *Letters in Applied Microbiology*, 68(2): 188-195. DOI: <https://doi.org/10.1111/lam.13106>
- Ammar AM, Mohamed AA, Abd El-Hamid MI, and El-Azzouny MM (2016). Virulence genotypes of clinical *Salmonella* serovars from broilers in Egypt. *Journal of Infection in Developing Countries*, 10(04): 337-346. DOI: <https://doi.org/10.3855/jidc.7437>
- Aslam M, Checkley S, Avery B, Chalmers G, Bohaychuk V, Gensler G, et al. (2012). Phenotypic and genetic characterization of antimicrobial resistance in *Salmonella* serovars isolated from retail meats in Alberta, Canada. *Food Microbiology*, 32(1): 110-117. DOI: <https://doi.org/10.1016/j.fm.2012.04.017>
- Awad A, Gwida M, Khalifa E, and Sadat A (2020). Phenotypes, antibacterial-resistant profile, and virulence-associated genes of *Salmonella* serovars isolated from retail chicken meat in Egypt. *Veterinary World*, 13(3): 440-445. DOI: <https://doi.org/10.14202%2Fvetworld.2020.440-445>

- Aziz SAA, Abdel-Latef GK, Shany SA, and Rouby SR (2018). Molecular detection of integron and antimicrobial resistance genes in multidrug resistant *Salmonella* isolated from poultry, calves and human in Beni-Suef governorate, Egypt. Beni-Suef University Journal of Basic and Applied Sciences, 7(4): 535-542. DOI: <https://doi.org/10.1016/j.bjbas.2018.06.005>
- Azizpour A (2018). Determining the antibiotic resistance patterns of isolated *Salmonella* from broiler flocks to 28 antimicrobial agents used in Iran. Journal of Comparative Pathobiology, 15(1): 2411-2420. Available at: [https://jcp.srbiau.ac.ir/article\\_11831.html?lang=en](https://jcp.srbiau.ac.ir/article_11831.html?lang=en)
- Badr H, Roshdy H, Sorour HK, AbdelRahman MAA, Erfan AM, Salem N, et al. (2021). Phenotypic and genotypic characterization of *Salmonella enterica* serovars isolated from imported poultry. Advances in Animals and Veterinary Sciences, 9(6): 823-834. DOI: <http://dx.doi.org/10.17582/journal.aavs/2021/9.6.823.834>
- Borges KA, Furian TQ, De Souza SN, Tondo EC, Streck AF, Salle CTP, et al. (2017). Spread of a major clone of *Salmonella enterica* serotype Enteritidis in poultry and in salmonellosis outbreaks in Southern Brazil. Journal of Food Protection, 80(1): 158-163. DOI: <https://doi.org/10.4315/0362-028x.jfp-16-299>
- Calayag AMB, Paclibare PAP, Santos PDM, Bautista CAC, and Rivera W (2017). Molecular characterization and antimicrobial resistance of *Salmonella enterica* from swine slaughtered in two different types of Philippine abattoir. Food Microbiology, 65: 51-56. DOI: <https://doi.org/10.1016/j.fm.2017.01.016>
- Campioni F, Bergamini AMM, and Falcão JP (2012). Genetic diversity, virulence genes and antimicrobial resistance of *Salmonella* Enteritidis isolated from food and humans over a 24-year period in Brazil. Food Microbiology, 32(2): 254-264. DOI: <https://doi.org/10.1016/j.fm.2012.06.008>
- Capuano F, Mancusi A, Capparelli R, Esposito S, and Proroga YT (2013). Characterization of drug resistance and virulotypes of *Salmonella* strains isolated from food and humans. Foodborne Pathogens and Disease, 10(11): 963-968. DOI: <https://doi.org/10.1089/fpd.2013.1511>
- Castro-Vargas RE, Herrera-Sánchez MP, Rodríguez-Hernández R, and Rondón-Barragán IS (2020). Antibiotic resistance in *Salmonella* spp. isolated from poultry: A global overview. Veterinary World, 13(10): 2070-2084. DOI: <https://doi.org/10.14202%2Fvetworld.2020.2070-2084>
- Cha SY, Kang M, Yoon RH, Park CK, Moon OK, and Jang HK (2013). Prevalence and antimicrobial susceptibility of *Salmonella* isolates in Pekin ducks from South Korea. Comparative Immunology, Microbiology and Infectious Diseases, 36(5): 473-479. DOI: <https://doi.org/10.1016/j.cimid.2013.03.004>
- Chuanchien R, and Padungtod P (2009). Antimicrobial resistance genes in *Salmonella enterica* isolates from poultry and swine in Thailand. Journal of Veterinary Medical Science, 71(10): 1349-1355. DOI: <https://doi.org/10.1292/jvms.001349>
- Diab MS, Zaki RS, Ibrahim NA, and Abd El Hafez MS (2019). Prevalence of multidrug resistance non-typhoidal *Salmonellae* isolated from layer farms and humans in Egypt. World Veterinary Journal, 9(4): 280-288. DOI: <https://dx.doi.org/10.36380/scil.2019.vwj35>
- Donado-Godoy P, Clavijo V, León M, Tafur MA, Gonzales S, Hume M, et al. (2012). Prevalence of *Salmonella* on retail broiler chicken meat carcasses in Colombia. Journal of Food Protection, 75(6): 1134-1138. DOI: <https://doi.org/10.4315/0362-028x.jfp-11-513>
- Doosti A, Mahmoudi E, Jami MS, and Mokhtari-Farsani A (2017). Prevalence of *aadA1*, *aadA2*, *aadB*, *strA* and *strB* genes and their associations with multidrug resistance phenotype in *Salmonella Typhimurium* isolated from poultry carcasses. The Thai Journal of Veterinary Medicine, 46(4): 691-697. DOI: <https://he01.tci-thaijo.org/index.php/tjvm/article/view/73827>
- Elhariri M, Elhelw R, Selim S, Ibrahim M, Hamza D, and Hamza E (2020). Virulence and antibiotic resistance patterns of extended-spectrum beta-lactamase-producing *Salmonella enterica* serovar Heidelberg isolated from broiler chickens and poultry workers: a potential hazard. Foodborne Pathogens and Disease, 17(6): 373-381. DOI: <https://doi.org/10.1089/fpd.2019.2719>
- Elkenany R, Elsayed MM, Zakaria AI, El-Sayed SAES, and Rizk MA (2019). Antimicrobial resistance profiles and virulence genotyping of *Salmonella enterica* serovars recovered from broiler chickens and chicken carcasses in Egypt. BMC Veterinary Research, 15: 124. DOI: <https://doi.org/10.1186/s12917-019-1867-z>
- Elsayed M, El-Gohary F, Zakaria A, and Gwida M (2019). Tracing of *Salmonella* contaminations throughout an integrated broiler production chain in Dakahlia governorate, Egypt. Pakistan Veterinary Journal, 39(4): 558-562. DOI: <https://doi.org/10.29261/pakvetj/2019.038>
- El-Sharkawy H, Tahoun A, El-Gohary AEGA, El-Abasy M, El-Khayat F, Gillespie T, et al. (2017). Epidemiological, molecular characterization and antibiotic resistance of *Salmonella enterica* serovars isolated from chicken farms in Egypt. Gut Pathogens, 9: 8. DOI: <https://doi.org/10.1186%2Fs13099-017-0157-1>
- Elshebrawy HA, Mahros MA, Abd-Elghany SM, Elgazzar MM, Hayashidani H, and Sallam KI (2021). Prevalence and molecular characterization of multidrug-resistant and  $\beta$ -lactamase producing *Salmonella enterica* serovars isolated from duck, pigeon, and quail carcasses in Mansoura, Egypt. LWT, 149: 111834. DOI: <https://doi.org/10.1016/j.lwt.2021.111834>
- ElSheikh M, Abdeen E, and Ammar A (2019). Molecular detection of some virulence genes of *Salmonella* serotypes isolated from poultry in Egypt. Journal of Current Veterinary Research, 1(1): 86-93. DOI: <https://doi.org/10.21608/jcvr.2019.36570>
- Firoozeh F, Zahraei-Salehi T, Shahcheraghi F, Karimi V, and Aslani MM (2012). Characterization of class I integrons among *Salmonella enterica* serovar Enteritidis isolated from humans and poultry. FEMS Immunology & Medical Microbiology, 64(2): 237-243. DOI: <https://doi.org/10.1111/j.1574-695X.2011.00883.x>
- Frye JG, and Jackson CR (2013). Genetic mechanisms of antimicrobial resistance identified in *Salmonella enterica*, *Escherichia coli*, and *Enterococcus* spp. isolated from US food animals. Frontiers in Microbiology, 4: 135. DOI: <https://doi.org/10.3389/fmicb.2013.00135>
- Ghetas A, Abdelbaki M, Fedawy H, Sedeeq DM, Bosila MA, Samy AA, et al. (2021). Antimicrobial resistance and virulence genotyping of different *Salmonella* serovars isolated from chickens in Egypt. Advances in Animal and Veterinary Sciences, 9(12): 2124-2131. DOI: <http://dx.doi.org/10.17582/journal.aavs/2021/9.12.2124.2131>
- Gopinath RS, Singh VP, Bhattacharya A, and Sharma B (1998). Molecular subtyping of the Indian isolates of *Salmonella gallinarum* using 16S rRNA gene loci PCR-Amplicon as a probe in Bangalore. Proceedings of the Second Pan Commonwealth Veterinary Conference, pp 893-899.
- Gyles CL (2008). Antimicrobial resistance in selected bacteria from poultry. Animal Health Research Reviews, 9(2): 149-158. DOI: <https://doi.org/10.1017/s1466252308001552>
- Habrun B, Šimpraga B, Kompes G, and Krstulović F (2012). Antimicrobial resistance and serotyping of *Salmonella enterica* subsp. *enterica* isolated from poultry in Croatia. Veterinarski Arhiv, 82(4): 371-381. Available at: <https://hrcak.srce.hr/file/125117>
- Hassan WH, Abed AH, Thabet A, and El Nady AM (2018). Genetic analysis of multidrug resistant *Salmonella* isolated from broiler chickens. Journal of Veterinary Medical Research, 25(1): 121-131. DOI: <https://doi.org/10.21608/jvmr.2018.43309>

- Hassan WH, Hassan HS, Hassan WM, Shany SA, and Osman GS (2021). Identification and characterization of *Salmonella* species isolated from broiler chickens. *Journal of Veterinary Medical Research*, 28(1): 21-29. DOI: <https://dx.doi.org/10.21608/JVMR.2021.89244.1041>
- Herrera-Sánchez MP, Rodríguez-Hernández R, and Rondón-Barragán IS (2020). Molecular characterization of antimicrobial resistance and enterobacterial repetitive intergenic consensus-PCR as a molecular typing tool for *Salmonella* spp. isolated from poultry and humans. *Veterinary World*, 13(9): 1771. DOI: <https://doi.org/10.14202/vetworld.2020.1771-1779>
- Huehn S, La Ragione RM, Anjum M, Saunders M, Woodward MJ, Bunge C, et al. (2010). Virulotyping and antimicrobial resistance typing of *Salmonella* enterica serovars relevant to human health in Europe. *Foodborne Pathogens and Disease*, 7(5): 523-535. DOI: <https://doi.org/10.1089/fpd.2009.0447>
- Ibrahim WA., Abd El-Ghany WA, Nasef SA, and Hatem ME (2014). A comparative study on the use of real time polymerase chain reaction (RT-PCR) and standard isolation techniques for the detection of *Salmonella* in broiler chicks. *International Journal of Veterinary Science and Medicine*, 2 (1): 67-71. DOI: <https://doi.org/10.1016/j.ijvsm.2013.11.001>
- Ibrahim S, Wei Hoong L, Lai Siong Y, Mustapha Z, CW Zalati CS, Aklilu E, et al. (2021). Prevalence of antimicrobial resistance (AMR) *Salmonella* spp. and *Escherichia coli* isolated from broilers in the East Coast of Peninsular Malaysia. *Antibiotics*, 10(5): 579. DOI: <https://doi.org/10.3390/antibiotics10050579>
- International Organization for Standardization (ISO) (2002). Microbiology of food and animal feeding stuffs-horizontal method for the detection of *Salmonella* spp., 4th edition. Geneva. Available at: <https://www.iso.org/obp/ui/#iso:std:iso:6579:ed-4:v1:en>
- Kapoor G, Saigal S, Elongavan A (2017). Action and resistance mechanisms of antibiotics: A guide for clinicians. *Journal of Anaesthesiology, Clinical Pharmacology*, 33(3): 300-305. DOI: [https://doi.org/10.4103%2Fjoacp.JOACP\\_349\\_15](https://doi.org/10.4103%2Fjoacp.JOACP_349_15)
- Karmi M (2013). Detection of virulence gene (*invA*) in *Salmonella* isolated from meat and poultry products. *International Journal of Genetics*, 3(2): 7-12. DOI: <https://doi.org/10.5829/idosi.ijg.2013.3.2.82204>
- Kauffman G (1974). Kauffmann White Scheme. *Acta Pathologica et Microbiologica Scandinavica*, 61: 385.
- Kuria KNJ (2023). Salmonellosis in Food and Companion Animals and Its Public Health Importance. *Salmonella - Past, Present and Future [Working Title]*. IntechOpen. <http://dx.doi.org/10.5772/intechopen.109324>
- Lamas A, Fernandez-No IC, Miranda JM, Vázquez B, Cepeda A, and Franco CM (2016). Prevalence, molecular characterization and antimicrobial resistance of *Salmonella* serovars isolated from northwestern Spanish broiler flocks (2011–2015). *Poultry Science*, 95(9): 2097-2105. DOI: <https://doi.org/10.3382/ps/pew150>
- Lee N, Kwon KY, Oh SK, Chang HJ, Chun HS, and Choi SW (2014). A multiplex PCR assay for simultaneous detection of *Escherichia coli* O157: H7, *Bacillus cereus*, *Vibrio parahaemolyticus*, *Salmonella* spp., *Listeria monocytogenes*, and *Staphylococcus aureus* in Korean ready-to-eat food. *Foodborne Pathogens and Disease*, 11(7): 574-580. DOI: <https://doi.org/10.1089/fpd.2013.1638>
- Li P, Zhang D, Li H, Pang J, Guo H, and Qiu J (2020). Establishment and application of multiplex PCR for simultaneously detecting *Escherichia coli*, *Salmonella*, *Klebsiella pneumoniae*, and *Staphylococcus aureus* in minks. *Frontiers in Veterinary Science*, 7: 588173. DOI: <https://doi.org/10.3389/fvets.2020.588173>
- Lu Y, Wu CM, Wu GJ, Zhao HY, He T, Cao XY, et al. (2011). Prevalence of antimicrobial resistance among *Salmonella* isolates from chicken in China. *Foodborne Pathogens and Disease*, 8(1): 45-53. DOI: <https://doi.org/10.1089/fpd.2010.0605>
- Ma M, Wang H, Yu Y, Zhang D, and Liu S (2007). Detection of antimicrobial resistance genes of pathogenic *Salmonella* from swine with DNA microarray. *Journal of Veterinary Diagnostic Investigation*, 19(2): 161-167. DOI: <https://doi.org/10.1177/104063870701900204>
- Maciorowski KG, Herrera P, Jones FT, Pillai SD, and Ricke SC (2006). Cultural and immunological detection methods for *Salmonella* spp. in animal feeds—a review. *Veterinary Research Communications*, 30(2): 127-137. DOI: <https://doi.org/10.1007/s11259-006-3221-8>
- Margot H, Stephan R, O'Mahony E, and Iversen C (2013). Comparison of rapid cultural methods for the detection of *Salmonella* species. *International Journal of Food Microbiology*, 163(2-3): 47-50. DOI: <https://doi.org/10.1016/j.ijfoodmicro.2013.02.013>
- Marshall BM, and Levy SB (2011). Food animals and antimicrobials: impacts on human health. *Clinical Microbiology Reviews*, 24(4): 718-733. DOI: <https://doi.org/10.1128/cmr.00002-11>
- Menghistu HT, Rathore R, Dhama K, and Agarwal RK (2011). Isolation, identification and polymerase chain reaction (PCR) detection of *Salmonella* species from field materials of poultry origin. *International Journal of Microbial Research*, 2(2): 135-142. Available at: [https://www.idosi.org/ijmr/ijmr2\(2\)11/7.pdf](https://www.idosi.org/ijmr/ijmr2(2)11/7.pdf)
- Mohamed M, Mohamed R, Gharieb R, Amin M, and Ahmed H (2021). Antimicrobial resistance, virulence associated genes and biofilm formation of *Salmonella* species isolated from different sources. *Zagazig Veterinary Journal*, 49(2): 208-221. DOI: <https://doi.org/10.21608/zvjz.2021.81722.1144>
- Mohamed MEM, and Suelam IIA (2010). Isolation of non-typhoid *Salmonella* from humans and camels with reference to its survival in abattoir effluents. *Global Veterinaria*, 5(6): 356-361. Available at: [https://www.idosi.org/gv/gv5\(6\)10/10.pdf](https://www.idosi.org/gv/gv5(6)10/10.pdf)
- Mottet, A., and Tempio, G. (2017). Global poultry production: Current state and future outlook and challenges. *World's Poultry Science Journal*, 73(2): 245-256. <https://doi.org/10.1017/S0043933917000071>
- Moussa IM, Gassem MA, Al-Doss AA, Sadik WAM, and Mawgood AA (2010). Using molecular techniques for rapid detection of *Salmonella* serovars in frozen chicken and chicken products collected from Riyadh, Saudi Arabia. *African Journal of Biotechnology*, 9(5): 612-619. DOI: <https://doi.org/10.5897/AJB09.1761>
- Murugkar HV, Rahman H, and Dutta PK (2003). Distribution of virulence genes in *Salmonella* serovars isolated from man & animals. *Indian Journal of Medical Research*, 117: 66-70. Available at: <https://pubmed.ncbi.nlm.nih.gov/12931840/>
- Nabil NM, and Yonis AE (2019). Isolation of *Salmonella* characterized by biofilm formation and disinfectant resistance from broiler chickens. *Alexandria Journal of Veterinary Sciences*, 62(2): 26-36. DOI: <http://dx.doi.org/10.5455/ajvs.57274>
- National Committee for Clinical Laboratory Standards (NCCLS) (2020). Performance standards for antimicrobial susceptibility testing: eleventh informational supplement. M100- S11. 30<sup>th</sup> edition, Wayne. PA. USA. Available at: [https://clsi.org/media/3481/m100ed30\\_sample.pdf](https://clsi.org/media/3481/m100ed30_sample.pdf)
- Oliveira SD, Rodenbusch CR, Ce MC, Rocha SLS, and Canal CW (2003). Evaluation of selective and non-selective enrichment PCR procedures for *Salmonella* detection. *Letters in Applied Microbiology*, 36(4): 217-221. DOI: <https://doi.org/10.1046/j.1472-765x.2003.01294.x>
- O'Regan E, McCabe E, Burgess C, McGuinness S, Barry T, Duffy G, et al. (2008). Development of a real-time multiplex PCR assay for the detection of multiple *Salmonella* serotypes in chicken samples. *BMC Microbiology*, 8: 165. DOI: <https://doi.org/10.1186/1471-2180-8-156>

- Osman KM, Marouf SH, Zolnikov TR, and AlAtfeehy N (2014). Isolation and characterization of *Salmonella enterica* in day-old ducklings in Egypt. *Pathogens and Global Health*, 108(1): 37-48. DOI: <https://doi.org/10.1179%2F2047773213Y.0000000118>
- Quinn PJ, Markey BK, Leonard FC, Hartigan P, Fanning S, and Fitzpatrick ES (2011). *Veterinary Microbiology and Microbial Disease*. 2<sup>nd</sup> Edition, Wiley-Blackwell Science. Available at: <https://www.wiley.com/enbe/Veterinary+Microbiology+and+Microbial+Disease%2C+2nd+Edition-p-9781405158237>
- Rabie NS, Khalifa NO, Radwan ME, and Afify JS (2012). Epidemiological and molecular studies of *Salmonella* isolates from chicken, chicken meat and human in Toukh, Egypt. *Global Veterinaria*, 8(2): 128-132. Available at: [https://fvtm.stafpu.bu.edu.eg/Zoonotic%20Diseases/957/publications/Nashwa%20Osman%20Khalefa\\_GLOBAL%20VET.pdf](https://fvtm.stafpu.bu.edu.eg/Zoonotic%20Diseases/957/publications/Nashwa%20Osman%20Khalefa_GLOBAL%20VET.pdf)
- Radwan IA, Abed AH, Abd Al-Wanis SA, Abd El-Aziz GG, and El-Shemy A (2016). Antibacterial effect of cinnamon and oreganium oils on multidrug resistant *Escherichia coli* and *Salmonellae* isolated from broiler chickens. *Journal of the Egyptian Veterinary Medical Association*, 76(2): 169-186.
- Raji MA, Kazeem HM, Magyigbe KA, Ahmed AO, Lawal DN, and Raufu IA (2021). *Salmonella* serovars, antibiotic resistance, and virulence factors isolated from intestinal content of slaughtered chickens and ready-to-eat chicken gizzards in the Ilorin Metropolis, Kwara State, Nigeria. *International Journal of Food Science*, 2021: 8872137. DOI: <https://doi.org/10.1155/2021/8872137>
- Ramatla TA, Mphuthi N, Ramaili T, Taioe MO, Thekisoe OM, and Syakalima M (2020). Molecular detection of virulence genes in *Salmonella* spp. isolated from chicken faeces in Mafikeng, South Africa. *Journal of the South African Veterinary Association*, 91(0): 1-7. DOI: <https://doi.org/10.4102%2Fjsava.v91i0.1994>
- Roca I, Akova M, Baquero F, Carlet J, Cavalieri M, Coenen S, et al. (2015). The global threat of antimicrobial resistance: science for intervention. *New Microbes and New Infections*, 6: 22-29. DOI: <https://doi.org/10.1016/j.nmni.2015.02.007>
- Sabry MA, Abdel-Moein KA, Abdel-Kader F, and Hamza E (2020). Extended-spectrum  $\beta$ -lactamase-producing *Salmonella* serovars among healthy and diseased chickens and their public health implication. *Journal of Global Antimicrobial Resistance*, 22: 742-748. DOI: <https://doi.org/10.1016/j.jgar.2020.06.019>
- Saleem G, Farooq U, Javed MT, Naseer R, Aslam HB, Mustafa G, et al. (2022). Pathobiological and immunohistochemical findings in broiler chickens naturally infected with *Salmonella enterica* serotype *Gallinarum* biotype *Gallinarum*. *Pakistan Veterinary Journal*, 42(1): 88-94. DOI: <https://doi.org/10.29261/pakvetj/2021.083>
- Samanta I, Joarder SN, Das PK, Sar TK, Bandyopadhyay S, Dutta TK, et al. (2014). Prevalence and antibiotic resistance profiles of *Salmonella* serotypes isolated from backyard poultry flocks in West Bengal, India. *Journal of Applied Poultry Research*, 23(3): 536-545. DOI: <https://doi.org/10.3382/japr.2013-00929>
- Shabnam M, and Kwai Lin T (2010). Isolation and molecular sub typing of *Salmonella enterica* from chicken, beef and street foods in Malaysia. *Scientific Research and Essays*, 5(18): 2713-2720. Available at: <https://eprints.um.edu.my/18919/>
- Shah DH, Zhou X, Addwebi T, Davis MA, Orfe L, Call DR, et al. (2011). Cell invasion of poultry-associated *Salmonella enterica* serovar Enteritidis isolates is associated with pathogenicity, motility and proteins secreted by the type III secretion system. *Microbiology*, 157(5): 1428-1445. DOI: <https://doi.org/10.1099/mic.0.044461-0>
- Shehata AA, Basiouni S, Elrazek AA, Sultan H, Tarabees R, Abd MS, et al. (2019). Characterization of *Salmonella enterica* isolated from poultry hatcheries and commercial broiler chickens. *Pakistan Veterinary Journal*, 39(4): 515-520. DOI: <https://doi.org/10.29261/pakvetj/2019.033>
- Siddique RA, Saxena M, and Lakhchaura BD (2009). PCR based rapid detection of *Salmonella* from poultry samples. *Indian Journal of Public Health*, 53(4): 226-228. Available at: <https://pubmed.ncbi.nlm.nih.gov/20469760/>
- Singh S, Yadav A, Singh S, and Bharti P (2010). Prevalence of *Salmonella* in chicken eggs collected from poultry farms and marketing channels and their antimicrobial resistance. *Food Research International*, 43(8): 2027-2030. DOI: <https://doi.org/10.1016/j.foodres.2010.06.001>
- World Health Organization (WHO) (2006). Progress report (2000-2005): building capacity for laboratory-based foodborne disease surveillance and outbreak detection and response. World Health Organization. DOI: <https://apps.who.int/iris/handle/10665/43424>
- Xu Y, Zhou X, Jiang Z, Qi Y, Ed-Dra A, Yue M (2020). Epidemiological investigation and antimicrobial resistance profiles of *Salmonella* isolated from breeder chicken hatcheries in Henan, China. *Frontiers in Cellular and Infection Microbiology*, 10: 497. DOI: <https://doi.org/10.3389/fcimb.2020.00497>
- Zou M, Keelara S, and Thakur S (2012). Molecular characterization of *Salmonella enterica* serotype Enteritidis isolates from humans by antimicrobial resistance, virulence genes, and pulsed-field gel electrophoresis. *Foodborne Pathogens and Disease*, 9(3): 232-238. DOI: <https://doi.org/10.1089/fpd.2011.1012>